

## Neonatal mucopolipidosis 2. The spontaneous evolution of early bone lesions and the effect of vitamin D treatment

### Report of two cases

U. E. Pazzaglia<sup>1</sup>, G. Beluffi<sup>2</sup>, C. Danesino<sup>3</sup>, P. V. Frediani<sup>4</sup>, G. Pagani<sup>5</sup> and G. Zatti<sup>1</sup>

<sup>1</sup> Clinica Ortopedica dell'Università di Pavia and <sup>2</sup> Servizio di Radiodiagnostica, IRCCS Policlinico S. Matteo, Pavia,

<sup>3</sup> Cattedra di Genetica Umana, Università di Sassari, Sassari, <sup>4</sup> Ospedale dei Bambini Umberto I, Brescia,

<sup>5</sup> Divisione di Patologia Neonatale, Ospedale S. Anna, Como, Italy

**Abstract.** Evolution of the early bone lesions in two children with mucopolipidosis 2 was followed from birth. The progression of the bone changes did not differ from healing of rickets. Low levels of 1,25-(OH)<sub>2</sub>-D<sub>3</sub> were found in one child and he was treated with vitamin D; resolution of the rachitic changes was more rapid than in the untreated child. It is suggested that in mucopolipidosis 2 bone reacts to two independent factors, one controlling calcium metabolism, the other depending on the primary lysosomal enzyme defect. Since ricket-like features are not present in the other mucopolipidoses or mucopolysaccharidoses, the defect of calcium metabolism seems to be related to the specific enzyme defect of mucopolipidosis 2.

When mucopolipidosis 2 is detected at birth peculiar radiographic changes of the bones have been described [1–5], which strongly resemble rickets or osteomalacia [6, 7]. Histologic examination confirmed the presence of rachitic lesions and signs of hyperparathyroidism [8]. At later ages the disease presents with the typical, Hurler-like signs of dysostosis multiplex. As neonatal mucopolipidosis 2 has an uneventful course, evolution of the radiographic changes has been documented in only a few cases [9, 10].

In the present study bone changes were followed from birth in two cases; on the basis of the previous histological findings [8] one child was treated with vitamin D and the results are described.


### Case reports

Patient 1 – G.L., male, was the 2300 g term product of a 36-year-old gravida-1 para-0 abortus-1 mother and a 38-year-old non consanguineous father. At birth Apgar scores were 8 and 9 at 1' and 5'

respectively. The child was noted to have wizened facial features, a long philtrum and gingival hypertrophy. The lower limbs were short with bowing of the tibiae.

Radiographic features are reported in the next section.

Piastrinopenia was present at birth, but by 12 days of age platelets count was normal. Karyogram was 46 XY; urinary MPS ex-



**Fig. 1.** Case 1. Evolution of radiographic changes in femura from 6 month to 1 year of age. No vitamin D or other treatment was given to the child

**Table 1.** Phospho-calcium metabolism parameters available in the two I-cell disease cases

|   | Case 1 |        |        | Case 2 |        |        |         |         |          |          |          |          |
|---|--------|--------|--------|--------|--------|--------|---------|---------|----------|----------|----------|----------|
| Age   | 3 days | 6 days | 1 year | 3 days | 5 days | 8 days | 20 days | 30 days | 2 months | 4 months | 6 months | 8 months |
| Calcium<br>Mg/dl  | 8.7    | 9.2    | 10.3   | 9.1    | 9.5    | 9.8    | 10.8    | 9.9     | 10.2     | 9.7      | 10.3     | 9.7      |
| Phosphorus<br>Mg/dl                                       | 5.1    | 3.4    | 4.85   | -      | 2.8    | 2.3    | 5.9     | 5       | 5.8      | 5.6      | 5.3      | 4.2      |
| Alcaline<br>phosphatase<br>mU/ml                          | 503    | 725    | 523    | -      | 2008   | -      | 6450    | 3796    | 2720     | 2055     | 1522     | 1243     |
| Calcium-urine<br>Mg/dl                                    | -      | -      | 5.07   | -      | -      | -      | 0.80    | -       | 9.26     | -        | -        | 13.4     |
| Phosphorus-urine<br>Mg/dl                                 | -      | -      | 88.5   | -      | -      | -      | 7.0     | -       | 49.4     | -        | -        | 102      |
| Creatinine-urine<br>Mg/dl                                 | -      | -      | -      | -      | -      | -      | -       | -       | -        | -        | -        | 38       |
| PTH<br>ng/ml (0.2-0.8) <sup>a</sup>                       | -      | -      | 0.39   | 1.6    | -      | -      | -       | -       | 1.16     | -        | -        | -        |
| 25-OH-D3<br>ng% (20-60) <sup>a</sup>                      | -      | -      | -      | -      | -      | -      | -       | > 80    | -        | -        | -        | -        |
| 1,25-(OH) <sub>2</sub> -D3<br>pg% (32.8 ± 7) <sup>a</sup> | -      | -      | -      | -      | -      | -      | -       | 26.1    | -        | -        | -        | -        |
| Idrossiprolin-urine<br>Mg/24 h (10-40) <sup>a</sup>       | -      | -      | -      | -      | -      | -      | 38.1    | -       | -        | -        | -        | -        |

<sup>a</sup> () Normal values**Table 2.** Acid hidrolases determinations in plasma and cultured fibroblasts

|             |                                | Case 1                          |                                 | Case 2                |                      |
|-------------|--------------------------------|---------------------------------|---------------------------------|-----------------------|----------------------|
|             |                                | 1 month                         | 1 year                          | birth                 | 6 month              |
| Plasma      | Beta-exosaminidase             | 457 (6.8 - 12.8) <sup>a,c</sup> | 12254.4 (500-3000) <sup>a</sup> | 13104.48 <sup>b</sup> | 9102.90 <sup>b</sup> |
|             | Alpha-mannosidase              | 58 (0.25- 0.59) <sup>a,c</sup>  | -                               | -                     | -                    |
|             | Alpha-fucosidase               | 45 (1.8 - 6.7) <sup>a,c</sup>   | -                               | -                     | -                    |
|             | Arylsulphatase A               | 32 (0.4 - 1) <sup>a,c</sup>     | 67.55 (0.5-2.5) <sup>a</sup>    | 39.77 <sup>b</sup>    | 73.05 <sup>b</sup>   |
|             | Alpha-n-acetil glucosaminidase | -                               | 259.59 (10-25) <sup>a</sup>     | 93.81 <sup>b</sup>    | 216.4 <sup>b</sup>   |
| Fibroblasts | Beta-exosaminidase             | -                               | 784.93 (2500-10000)             | 948.37 <sup>b</sup>   | -                    |
|             | Alpha-fucosidase               | -                               | 3.16 (30-130) <sup>a</sup>      | 2.52 <sup>b</sup>     | -                    |
|             | Arylsulphatase A               | -                               | 12.81 (150-500) <sup>a</sup>    | 20.61 <sup>b</sup>    | -                    |
|             | Beta-galattosidase             | -                               | 12.73 (300-800) <sup>a</sup>    | 18.73 <sup>b</sup>    | -                    |

Activity expressed as nm/mg/hour in fibroblasts, as nm/ml/hour in serum

<sup>a</sup> () Normal values<sup>b</sup> Normal values as in the second column<sup>c</sup> Normal values are different in case 1 at 1 month because determinations were carried out in another lab

cretion was normal. Mucopolipidosis 2 was diagnosed at two months of age on the basis of the increased activity of beta-hexosaminidase, alfa-mannosidase, arylsulfatase A and alfa-L-fucosidase in serum (Table 1).

Growth was slow and motor development delayed. There was progressive coarsening of the facial features and gingival hypertrophy, with fully expressed Hurler-like features by six months of age.

Patient 2 - C.D., male, was the third son of healthy non-consanguineous parents. The older sister was normal but a brother with mucopolipidosis 2 died at 1 year of age. Genetic counselling was given to the parents after the second pregnancy, but they refused prenatal diagnosis. Fetal bone dysplasia was documented by ultrasound performed in the 6th month of pregnancy. The child was delivered at 38 week by Caesarean section.

Weight was 2270 g (below the 3rd percentile); with an Apgar score of 8 at birth. The patient had a wizened facies, high forehead, flat nose with anteverted nostrils, high-arched palate, long philtrum and gingival hypertrophy. Length was 42.5 cm (below the 3rd percentile) and head circumference 32 cm (15th percentile).

A deformed thorax, splenomegaly, short upper limbs and arachnodactyly, short lower limbs, club feet and striking radiographic skeletal abnormalities were present.

The diagnosis of mucopolipidosis 2 was established soon after birth on the basis of increased serum hydrolase activity, inclusions in cultured fibroblasts and decreased hydrolase activity in leukocytes and cultured fibroblasts (Table 1). Serum calcium, phosphorus, alkaline phosphatase, parathormone and 1,25-(OH)<sub>2</sub>-D3 as well as urine calcium and phosphorus, were determined and are reported in Table 2.

Treatment with 25-OH-D<sub>3</sub> (5 µg/Kg) was begun at 7 days of age; the dose was increased on the 15th (10 µg/Kg) and 23rd day (15 µg/Kg) and maintained up to 6 months of age.



**Fig. 2.** Case 1. Age 6 month. Stippling of tarsal bones. Cupping of the distal metaphyses of tibia and fibula is still evident as well as mesh-like trabecular structure of the bone

### Radiographic findings


Severe osteopenia was present in case 1 at birth. A thoraco-lumbar kyphosis of the spine was present. The lower thoracic and lumbar vertebrae had an oval shape. Thorax was narrow and the ribs were wide. The long bones had a mesh-like trabecular structure with periosteal cloaking, metaphyseal cupping and bowing of the tibiae and femora (with a sharp angle of the left femur). Stippling of the tarsal bones was present.

Although calcification of the long bones had improved at six months of age they were still osteoporotic. A transverse, radiopaque band was present in the proximal humeral and both femoral metaphyses. Periosteal calcifications appeared more dense; in the humeri they extended for the entire length of the diaphysis and doubled the thickness and size of the bone; in the femora it was restricted to the proximal part, where it filled the outer space between the transverse, radiopaque band and the diaphysis (Fig. 1). Cupping of the distal tibial and fibular metaphyses was still evident (Fig. 2). At 12 months bone density was normal and morphology also approached a normal pattern with well structured cortical bone in the diaphyses and well calcified trabeculae in the metaphyses. Bowing was still evident. The humeri had undergone remodelling and were shorter than normal but they were not as thick as before and the major metaphyseal abnormalities had disappeared. A similar evolution was observed in the femora (Fig. 1).

On the contrary, the classic Hurler-like signs of expanded and proximally pointed metacarpals and bullet-shaped phalanges had become apparent in the hands.



**Fig. 3.** Case 2. Evolution of radiographic changes in the left humerus from birth to 6 months of age. 25-OH-D<sub>3</sub> was given to the child from the 7th day



**Fig. 4.** Case 2. Right lower limb changes from birth to 6 month of age



**Fig. 5.** Case 2. X-rays of the hand at 6 month of age. Initial pointing of metacarpals and bullet shape of phalanges is observed

Case 2 showed the similar deformities of the spine and thorax as well as marked osteopenia and structural abnormalities of the long bones. The changes in the radiographic features (humerus and femur) from birth to six months of age were qualitatively similar to those observed in case 1, but with a much more rapid evolution illustrated in Figs. 3 and 4; this process proceeded through appearance of the metaphyseal, radiopaque band, increased

radiopacity of bone and subperiosteal apposition, remodeling to the funnelled shape of the metaphyses. Progression was the same in the upper and lower limb, but at each control the femur was in a more advanced phase than the humerus. At six months initial pointing of metacarpals and coning of phalanges was observed (Fig. 5).

## Discussion

No satisfactory explanation of the difference in the skeletal abnormalities observed in mucopolipidosis 2 at birth and dysostosis multiplex few months later has been presented to now. Autopsy findings in two cases showed ricket-like lesions of the bones and hyperparathyroidism [8], giving a pathological basis to the radiographic findings in neonatal cases.

This study further supports such a hypothesis since low levels of  $1,25-(\text{OH})_2\text{-D}_3$  were demonstrated in one child. The spontaneous evolution of early mucopolipidosis 2 lesions in the other did not differ from healing of rickets. On this assumption case 2 was treated from birth with vitamin D with faster resolution of the rachitic signs.

The tendency to spontaneous resolution of rachitic signs and at the same time the progression to lesions characteristic of a storage disease could suggest a response of bone to two independent factors, one controlling calcium metabolism and the other related to the primary lysosomal enzyme defect. This hypothesis raises two questions:

- 1) The transient nature of the defective calcium controlling mechanism.
- 2) Relationship to the primary enzyme defect.

As far as the first point is concerned, the observation of low levels of 1,25-(OH)<sub>2</sub>-D<sub>3</sub> in case 2 suggests a defective hydroxylation of vitamin D metabolites during intrauterine life and the spontaneous recovery after birth suggests subsequent correction of the defect.

As far as the second point is concerned the neonatal, transient, rickets-like changes, to the best of our knowledge, has never been reported in any other storage disease. The specific defect of Mucopolipidosis 2 is the lack of a phospho-transferase and a reduced cellular uptake of many lysosomal enzymes; since this defect is present during the fetal life and continues unchanged in the post-natal life, it does not seem causally related to the calcium metabolism defect, but could suggest a yet unexplained relationship with vitamin D metabolism in the "maternal environment".

## References

1. Cipolloni C, Boldrini A, Donti E, Coppa GV (1980) Neonatal mucopolipidosis II (I-cell disease): clinical, radiological and biochemical studies in a case. *Helv Paediatr Acta* 35: 85
2. Iannacone G, Capotorti L (1969) Contribution au syndrome dit "Pseudo-Hurler". Observations de deux soeurs avec altérations osseuses particulièrement sévères. *Ann Radiol (Paris)* 12: 355
3. Joannard A, Bost M, Pont J, Dieterlen M, Frappat P, Beau-doing A (1976) La mucopolipidose type II. Etude de deux observations familiales. Aspect cliniques et biochimiques. *Pediatrie* 29: 825
4. Lemaitre L, Remy T, Farriaux JP, Dhondt JL, Walbaum R (1978) Radiological signes of mucopolipidosis II or I-cell disease. *Pediatr Radiol* 7: 97
5. Patriquin HB, Kaplan P, Kind HO, Giedion A (1977) Neonatal mucopolipidosis II (I-cell disease): clinical and radiological features in three cases. *AJR* 129: 37
6. Maroteaux P (1971) Les mucopolipidoses. *Jourees Parisienne Pediatrie* Flanmanis N, Paris, p357
7. Maroteaux P, Hors-Cayla MC, Pont G (1970) La Mucopolipidose de type II. *Presse Med* 78: 179
8. Pazzaglia UE, Beluffi G, Campbell JB, Bianchi E, Colavita N, Diard F, Gugliantini P, Hirche U, Kozlowski K, Marchi A, Nayanar V, Pagani G (in press) Mucopolipidosis II: Correlation between radiological features and histopathology of the bones. *Pediatr Radiol* 19: 406
9. Leroy JG, Spranger JW, Feingold ME, Opitz JM, Crocker AC (1971) I-cell disease. A clinical picture. *J Pediatr* 79: 360
10. Spritz RA, Doughty RA, Spackman TJ, Murnane MT, Coates PM, Koldovsky O, Zackai EH (1978) Neonatal presentation of I-cell disease. *Pediatr* 93: 954

Received: 1 December 1988; accepted: 27 January 1989

Prof. U.E. Pazzaglia  
Clinica Ortopedica dell'Università di Pavia  
IRCCS S.Matteo  
Via Taramelli 3  
I-27100 Pavia  
Italy