


## Regular Article

# Prenatal maternal stress during the COVID-19 pandemic and infant regulatory capacity at 3 months: A longitudinal study

Livio Provenzi<sup>1</sup> , Serena Grumi<sup>1</sup>, Lilia Altieri<sup>2</sup>, Giulia Bensi<sup>3</sup>, Emanuela Bertazzoli<sup>4</sup>, Giacomo Biasucci<sup>3</sup>, Anna Cavallini<sup>5</sup>, Lidia Decembrino<sup>2</sup>, Rossana Falcone<sup>2</sup>, Anna Freddi<sup>6</sup>, Barbara Gardella<sup>7</sup>, Roberta Giacchero<sup>4</sup>, Roberto Giorda<sup>8</sup>, Elena Grossi<sup>3</sup>, Paola Guerini<sup>2</sup>, Maria Luisa Magnani<sup>2</sup>, Paola Martelli<sup>9</sup>, Mario Motta<sup>9</sup>, Renata Nacinovich<sup>5,10</sup>, Dario Pantaleo<sup>2</sup>, Camilla Pisoni<sup>7</sup>, Federico Prefumo<sup>9,11</sup>, Laura Riva<sup>6</sup>, Barbara Scelsa<sup>6</sup>, Maria V. Spartà<sup>4</sup>, Arsenio Spinillo<sup>7,12</sup>, Patrizia Vergani<sup>5,13</sup>, Simona Orcesi<sup>1,12</sup>, Renato Borgatti<sup>1,12</sup> and MOM-COPE Study Group†

<sup>1</sup>Child Neurology and Psychiatry Unit, IRCCS Mondino Foundation, Pavia, Italy; <sup>2</sup>ASST Pavia, Pavia, Italy; <sup>3</sup>Guglielmo da Saliceto Hospital, Piacenza, Italy; <sup>4</sup>ASST Lodi, Lodi, Italy; <sup>5</sup>San Gerardo Hospital, Monza, Italy; <sup>6</sup>ASST Sacco Fatebenefratelli, Milano, Italy; <sup>7</sup>Fondazione IRCCS Policlinico San Matteo, Pavia, Italy; <sup>8</sup>Scientific Institute IRCCS E. Medea, Bosisio Parini, Italy; <sup>9</sup>ASST Spedali Civili, Brescia, Italy; <sup>10</sup>School of Medicine and Surgery and Milan Center for Neuroscience, University of Milano Bicocca, Milano, Italy; <sup>11</sup>Department of Obstetrics and Gynaecology, University of Brescia, Brescia, Italy; <sup>12</sup>Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy and <sup>13</sup>Fondazione MBBM, Monza, Italy

## Abstract

The COVID-19 pandemic is a global traumatic experience for citizens, especially during sensitive time windows of heightened plasticity such as pregnancy and neonatal life. Pandemic-related stress experienced by mothers during pregnancy may act as an early risk factor for infants' regulatory capacity development by altering maternal psychosocial well-being (e.g., increased anxiety, reduced social support) and caregiving environment (e.g., greater parenting stress, impaired mother–infant bonding). The aim of the present longitudinal study was to assess the consequences of pandemic-related prenatal stress on infants' regulatory capacity. A sample of 163 mother–infant dyads was enrolled at eight maternity units in northern Italy. They provided complete data about prenatal stress, perceived social support, postnatal anxiety symptoms, parenting stress, mother–infant bonding, and infants' regulatory capacity at 3 months of age. Women who experienced emotional stress and received partial social support during pregnancy reported higher anxious symptoms. Moreover, maternal postnatal anxiety was indirectly linked to the infants' regulatory capacity at 3 months, mediated by parenting stress and mother–infant bonding. Dedicated preventive interventions should be delivered to mothers and should be focused on protecting the mother–infant dyad from the detrimental effects of pandemic-related stress during the COVID-19 healthcare emergency.

**Keywords:** anxiety, COVID-19, epidemic, maternal bonding, prenatal stress, regulatory capacity, social support, temperament

(Received 9 March 2021; revised 3 June 2021; accepted 4 June 2021)

## Introduction

During the first months of 2020, Italy was dramatically affected by the outbreak of the coronavirus disease of 2019 (COVID-19). The rapid spread and high contagion rate of COVID-19 together with the lockdown resulting from containment strategies soon contributed to an emerging scenario of global stress and a threat to public

health (Barello, Palamenghi, & Graffigna, 2020; Xiong et al., 2020). Exposure to stress may be especially detrimental during pregnancy, which is an early sensitive period characterized by high plasticity and heightened susceptibility to adverse environmental conditions (Davis & Narayan, 2020). In this paper, we document how early exposure to pandemic-related stress during pregnancy may affect infants' behavioral development of regulatory skills through a complex pathway that involves several factors related to reduced maternal psychosocial well-being and altered caregiving environment.

Recent studies suggested that pregnant women may not be at high risk for severe COVID-19 illness, as most mothers who were positive for the virus have been discharged without major health complications (Breslin et al., 2020; Schwartz, 2020; Zaigham & Andersson, 2020). Notwithstanding, even in the absence of critical COVID-19 clinical conditions, the exposure to an unprecedented pandemic may still result in heightened levels of stress for women during pregnancy and this may increase the risk for affective problems, such as anxiety symptomatology.

**Author for Correspondence:** Livio Provenzi, Child Neurology and Psychiatry Unit, IRCCS Mondino Foundation, via Mondino 2, 27100 Pavia, Italy. E-mail: [livio.provenzi@mondino.it](mailto:livio.provenzi@mondino.it)

†Additional members of the MOM-COPE project: Elisa Bettiga<sup>3</sup>, Emma Bonetti<sup>9</sup>, Renza Bonini<sup>3</sup>, Elisa Cavaleri<sup>9</sup>, Giovanna Centinaio<sup>4</sup>, Andrea Citterio<sup>8</sup>, Giuliana Del Campo<sup>2</sup>, Mattia Dominioni<sup>7</sup>, Andrea Gritti<sup>7</sup>, Gaia Kullmann<sup>5</sup>, Laura Malerba<sup>9</sup>, Eloisa Mariani<sup>5,13</sup>, Fabiana Mambretti<sup>8</sup>, Cristiana Pavesi<sup>3</sup>, Benedetta C. Pietra<sup>4</sup>, Caterina Sabatini<sup>4</sup>, Pierangelo Veggiotti<sup>6</sup>, Maria Luisa Ventura<sup>5,13</sup>, Marco Villa<sup>8</sup>, Sonia Zatti<sup>9</sup>, Marzo Zecca<sup>7</sup>

**Cite this article:** Provenzi L et al (2021). Prenatal maternal stress during the COVID-19 pandemic and infant regulatory capacity at 3 months: A longitudinal study. *Development and Psychopathology* 1–9. <https://doi.org/10.1017/S0954579421000766>

Recent studies reported high levels of stress and reduced psychosocial well-being among pregnant women during the pandemic (López-Morales *et al.*, 2021; Pope, Olander, Leitao, Meaney, & Matvienko-Sikar, 2021). Anxiety was among the most reported psychological symptoms in pregnant women and mothers in different countries hit by the COVID-19 pandemic (Cameron *et al.*, 2020; Lebel, MacKinnon, Bagshawe, Tomfohr-Madsen, & Giesbrecht, 2020; Racine *et al.*, 2021; Salehi, Rahimzadeh, Molaei, Zaheri, & Esmaelzadeh-Saeieh, 2020). During the healthcare emergency, women disclosed feelings of being unprepared to deliver (Preis, Mahaffey, Heiselman, & Lobel, 2020). Of note, many of these studies were cross-sectional and reported on maternal anxiety or stress related to the pandemic during pregnancy, but not in the postpartum period. As postnatal anxiety may significantly affect caregiving behaviors and infants' behavioral development (Field, 2018), the effects of pandemic-related stress on postnatal maternal anxious symptoms are warranted to be specifically investigated.

Moreover, because of COVID-19 mitigation strategies, mothers may have experienced reduced social support during pregnancy and this may have in turn contributed to further elevate their levels of distress and anxiety (Lebel *et al.*, 2020). The relative absence of fathers during and after delivery may have further contributed to maternal anxiety, as documented in previous research conducted during the COVID-19 pandemic (Lista & Bresesti, 2020). It is well known that perceived social support during pregnancy may result in a protective buffering effect in the face of prenatal stress on the subsequent risk of maternal anxiety, even in the case of traumatic events (Morikawa *et al.*, 2015; Tani & Castagna, 2017; Xie *et al.*, 2010; Xie, He, Koszycki, Walker, & Wen, 2009). For example, in a large longitudinal cohort, social support from the partner and significant others predicted significant decreases in stress and anxiety in pregnant women (Racine, Plamondon, Hentges, Tough, & Madigan, 2019). Similarly, the Iowa flood study (Brock *et al.*, 2015) demonstrated that social support might protect women from developing severe anxious symptomatology after exposure to traumatic stress events.

Extensive literature also suggested that prenatal stress and postpartum maternal anxiety might constitute relevant risk factors for maternal feelings of emotional closeness to the newborn, usually referred to with the term bonding (Field, 2018; Matvienko-Sikar, Murphy, & Murphy, 2018; Obrochta, Chambers, & Bandoli, 2020). Maternal bonding represents the emotional attachment between a caregiver and her infant; it develops shortly after birth and underpins several dimensions of parental caregiving skills, including sensitivity, pleasure for the interaction, and closeness (Van Bussel, Spitz, & Demyttenaere, 2010). Of note, maternal bonding may be challenged or impaired in the presence of prenatal stress, health-related risk, and postnatal anxiety (Fallon, Silverio, Halford, Bennett, & Harrold, 2021; Nicol-Harper, Harvey, & Stein, 2007; Provenzi *et al.*, 2017). For example, high levels of anxiety may be significantly associated with reduced maternal bonding at 2 weeks after delivery (Daglar & Nur, 2018) and at 4 months (Tietz, Zietlow, & Reck, 2014).

Prenatal maternal stress is also an adverse early experience that may affect a wide range of behavioral, emotional, and cognitive outcomes in infants and children (Su *et al.*, 2015; Zhu *et al.*, 2014), including infants' regulatory capacity during the first months of life (Class *et al.*, 2014; Van den Bergh *et al.*, 2020). The infant's regulatory capacity is a central component of temperament (Posner & Rothbart, 2000) that is highly susceptible to the

alterations of maternal psychosocial well-being and early caregiving environment (Gartstein & Skinner, 2018; Gunning, Halligan, & Murray, 2013). Prenatal stress has been previously found to be significantly associated with infants' temperament and regulatory skills in 3-month-old infants (Huizink, Robles De Medina, Mulder, Visser, & Buitelaar, 2002; Lin, Crnic, Luecken, & Gonzales, 2014) as well as in older children (Gutteling *et al.*, 2005). Similarly, maternal self-reported stress during pregnancy and postnatal anxiety have been especially linked with decreased infants' regulatory capacity, controlling for potential confounders (*i.e.*, birth weight, maternal postnatal well-being, and psychosocial risks) (Fuller, Messito, Mendelsohn, Oyeku, & Gross, 2018; Gutteling *et al.*, 2005; Huizink *et al.*, 2002; Lin *et al.*, 2014).

Maternal prenatal stress may affect infants' early temperament and regulatory capacity also indirectly, through a set of variables related to mothers' psychosocial well-being and altered caregiving environment. For example, in 5-month-old infants, maternal bonding was found to mediate the association between maternal anxiety and infants' regulatory capacities (Müller *et al.*, 2016), further suggesting that the link between maternal stress or affective problems and infants' temperament may be at least partially mediated by postpartum caregiving dimensions. Parenting stress is another variable that contributes to define the caregiving environment and that may mediate the association between maternal stress and temperament. For example, Sheinkopf *et al.* (2006) have reported in a large cohort of at-risk infants exposed to prenatal stress that the relation between neonatal characteristics and temperament might be moderated by maternal parenting stress.

Moreover, the presence of early problems in infants' regulatory capacity may be a risk factor for later socioemotional and behavioral issues during childhood (Feldman, 2015; Wittig & Rodriguez, 2019). For instance, infants with low regulatory capacity were found to be at greater risk for internalizing and externalizing behavioral problems during preschool age (Gartstein, Putnam, & Rothbart, 2012). Similarly, deficits in the attentive dimensions of infants' regulatory skills were linked with increased internalizing behaviors in children aged 6–10 years (Eisenberg *et al.*, 2009). As such, the early identification of regulatory capacity difficulties is key to promote preventive interventions, and the potential negative effects of prenatal maternal stress during the COVID-19 pandemic on infants' temperamental traits of regulatory capacity should not be underestimated. A longitudinal and prospective assessment of the psychosocial and caregiving pathways linking pandemic-related stress during pregnancy and infants' behavioral outcomes is warranted.

### *The present study*

Although extensive research provided evidence of the short-term and long-lasting effects of maternal prenatal stress on infants' developmental outcomes, there is a paucity of studies on the effects of pandemic-related prenatal stress on infants' temperament. As the present COVID-19 healthcare emergency should be considered as a global traumatic experience (Masiero, Mazzocco, Harnois, Cropley, & Pravettoni, 2020; Provenzi & Tronick, 2020), efforts should be dedicated to evaluating the potential indirect effects of pandemic-related prenatal stress on infants' development. The present healthcare emergency is a quasi-experimental condition suitable to study prospectively the short- and long-term effects of prenatal maternal stress in a community sample of low-risk women. Previous literature provides examples of quasi-experimental research investigating the effects

of large-scale disasters on mother–infant health. For example, studies conducted on the effects of the ice storm in Canada (Laplante, Brunet, & King, 2016) or the 2011 Australian floods (Simcock et al., 2017) provided evidence of the impact of prenatal stress on infant's outcomes, including temperament.

In order to understand the indirect impact of the pandemic-related prenatal stress on the development of infants' regulatory capacity, we launched a multi-centric, prospective longitudinal study involving multiple maternity units in northern Italy: measuring the outcomes of maternal COVID-19-related prenatal exposure (MOM-COPE) project (Provenzi et al., 2020c). This research project features multiple assessments of maternal mental health as well as infants' developmental outcomes from birth to age 12 months and integrates self-report, behavioral, and epigenetic measures. In the present study, we report on the effects of pandemic-related prenatal stress on infants' temperament at 3 months. We aimed to assess the relationship between maternal pandemic-related prenatal stress and the development of regulatory capacity in infants at 3 months. We hypothesized that higher prenatal stress would result in reduced regulatory skills in infants. Nonetheless, according to the aforementioned literature, we also tested specific mediators of this relationship related to maternal psychosocial status (i.e., postnatal anxiety, social support) and caregiving environment (parenting stress, maternal bonding).

## Method

### Participants and procedures

This study is part of the longitudinal MOM-COPE research project. The fully detailed description of this project is reported elsewhere (Provenzi et al., 2020c). Here we report on a sample of 163 mothers enrolled from May 2020, who provided complete data for prenatal ( $T_0$ ), neonatal ( $T_1$ ), and 3-month ( $T_2$ ) assessments by January 2021 (Figure 1). The attrition rate between  $T_0$  and  $T_2$  among enrolled mothers whose infants had reached 3 months of age by January 2021 was 23.1%. Mothers were included if at least 18 years old, in the absence of prenatal and perinatal diseases or injuries, if they delivered at term (i.e., from 37 + 0 to 41 + 6 weeks of gestation), and if they tested negative for SARS-CoV-2 at delivery. Mothers were first contacted at antepartum classes or immediately following the postpartum period. Sociodemographic and neonatal data were obtained from medical records. Within 48 hr from delivery, the mothers filled in a set of questionnaires to provide retrospective quantitative measures of prenatal COVID-19-related stress and perceived social support, as well as current postnatal anxious symptoms. When infants were approaching the age of 3 months, mothers received a second e-mail with the request to fill in additional questionnaires on postnatal mother–infant bonding, parenting stress, and infants' regulatory capacities. The study was approved by the Ethics Committees of the IRCCS Mondino Foundation (Pavia, Italy) and the participating hospitals. All mothers provided informed consent to participate in the study.

### Measures

#### Prenatal measures, $T_0$

Maternal sociodemographic information included age, educational level, and occupational status. At delivery, mothers retrospectively reported on their physical exposure to the virus and

prenatal COVID-19-related stress during the last trimester of pregnancy through ad hoc questionnaires (see Table S1 in the Supplementary Material). The direct (own positivity with or without symptoms) or indirect (positivity, hospitalization or death of relatives or significant others) physical exposure to COVID-19 was assessed with seven dichotomous items (0, *no*; 1, *yes*). The score was obtained by computing the sum of each item's score (range 0–7). The level of pandemic-related stress was assessed with six 5-point Likert scale items on the emotional stress response to the COVID-19 emergency (see Table S1 in the Supplementary Material) and rated from 1 (not at all) to 5 (very much). The pandemic-related stress score was obtained by computing a mean of the ratings obtained for each item (raw score range: 1–5). Prenatal social support was assessed and quantified using the global score of the Multidimensional Scale of Perceived Social Support, MSPSS (Zimet, Dahlem, Zimet, & Farley, 1988).

#### Neonatal measures, $T_1$


Neonatal characteristics (i.e., sex, gestational age, birth weight, Apgar at minute 5, and mode of delivery) were collected from medical records. Postnatal maternal state anxiety was assessed with the well-validated State-Trait Anxiety Inventory, STAI-Y (Spielberger, Gorsuch, & Luschene, 1983). The raw score ranged between 20 and 80, with 40 as a reliable cut-off score for clinical risk.

#### Postnatal measures, $T_2$

At infants' age of 3 months, mother–infant bonding was assessed with the Maternal Postpartum Attachment Scale, MPAS (Condon & Corkindale, 1998). It consists of 19 statements rated on a 2-, 4-, or 5-point scale response option. All responses were recoded to represent a score of 1 (low attachment) to 5 (high attachment) to ensure equal weight for all questions. The MPAS total score ranged from 19 to 95 with low scores indicating a problematic mother-to-infant bond. Maternal parenting stress was measured using the Parenting Stress Index Short Form, PSI-SF (Abidin, 1995). It includes 36 items that provide a global score of parental stress and three 0 subscales scores: parental distress, difficult child, and parent–child dysfunctional interactions. Finally, infants' regulatory capacity was assessed using the homonymous factor of the short-form version of the infant behavior questionnaire – revised, IBQ-R (Gartstein & Rothbart, 2003). The IBQ-R items are rated on a 7-point scale. The regulatory capacity includes items that index the following dimensions of infants' temperament: cuddliness, orienting, low-intensity pleasure, and soothability.

#### Plan of analysis

Potential associations among demographic/clinical characteristics and the variables of interest were explored through Pearson's bivariate correlations. Mothers with at least one direct or indirect exposure to COVID-19 were compared to counterparts with no exposure by means of independent-sample *t* tests. The path analysis model focused on the effects of maternal psychosocial well-being (i.e., *prenatal stress* and *prenatal support*), postnatal *maternal anxiety*, and caregiving environment variables (i.e., *mother–infant bonding* and *parenting stress*) on infants' *regulatory capacity* at 3 months. The final model included all the direct and indirect effects resulting in 17 free parameters and four degrees of freedom. Parameters were estimated using the maximum likelihood method. The covariances between *prenatal stress* and *prenatal support* (prenatal assessment) as well as between *parenting stress* and *mother–infant bonding* (3-month assessment) were



**Figure 1.** Schematic overview of the prenatal, perinatal, and 3-month assessments included in the study design. Note: prenatal measures were retrospectively assessed at childbirth. Perinatal data (T<sub>1</sub>) obtained 12–48 hr from delivery.

**Table 1.** Descriptive statistics of participants and study variables

		N	%
Delivery	Vaginal	109	66.9
	Elective cesarean	29	17.8
	Operative	13	8.0
	Emergency cesarean	12	7.3
Infants' sex	Females	83	50.9
	Males	80	49.1
Mothers' occupational status	Employed	146	89.6
	Unemployed	17	10.4
Fathers' occupational status	Employed	161	98.8
	Unemployed	2	1.2
Fathers' access to delivery	Yes	141	86.5
	No	22	13.5
Fathers' access during hospitalization	Yes	50	30.7
	No	113	69.3
	Mean	SD	Range
Mothers' age	33.79	4.83	18; 47
Mothers' educational level (years of study)	14.96	3.26	8; 22
Fathers' educational level (years of study)	13.26	3.37	8; 22
Infants' gestational age (weeks)	39.7	1.05	37; 42
Infants' birthweight (kg)	3.34	0.39	2.43; 4.34
Infants' Apgar (minute 5)	9.23	0.57	8; 10
Prenatal COVID-19-related maternal stress	2.49	0.67	1; 4.17
Prenatal maternal social support	6.06	1.09	1.6; 7
Postnatal maternal state anxiety	34.56	10.15	20; 74
Postnatal maternal parenting stress	56.87	15.77	36; 112
Postnatal mother–infant bonding	82.03	5.84	65.2; 93
Infant regulatory capacity	5.20	0.71	3.39; 6.8

also included in the model. This model was built according to the literature reviewed and reported in the introduction section of this manuscript and the final set of effects included in the path analysis was selected to maximize parsimony and model fit. A detailed description of the approximation models that led to the final path

analysis model tested and their relative indexes of fit is reported in the Supplementary Material, Supplementary File S1. All analyses were conducted using R (R Core Team, 2020) and the path analysis was performed using the *lavaan* package (Rosseel, 2012). The following indexes were used to confirm the goodness of fit of the

model: nonsignificant chi-square statistic, comparative fit index (CFI) and Tucker–Lewis index (TLI) close to .95, root mean squared error of approximation (RMSEA) smaller than .06, root mean square residual (SRMR) smaller than .08.

## Results

The characteristics of the present sample are summarized in Table 1. Subjects with complete data and those who were excluded from the analyses for missing data points between  $T_0$  and  $T_2$  did not show any statistically significant difference in sociodemographic characteristics and in the variables of interest. For what pertains, the exposure to COVID-19 (see Table 2), no mothers reported to have tested positive for SARS-CoV-2 during pregnancy. Nonetheless, up to one third of the sample was living in the first hotspot area of COVID-19 outbreak in northern Italy. Twenty-four percent of the sample ( $n = 39$ ) had at least one relative or close friend that needed intensive care in the hospital due to COVID-19 and 15% of the women ( $n = 24$ ) experienced the loss of a loved one. No gender-related statistically significant differences emerged for sociodemographic characteristics and variables of interest.

Maternal anxiety symptoms were above the clinical cut-off in 41 mothers (25.2%). The findings of the preliminary correlations are reported in Table 3. No significant associations emerged for demographic and clinical characteristics. Similarly, no differences in stress and anxiety emerged between mothers with and without at least one direct or indirect physical exposure to COVID-19. As such, these were not included in the path analysis model.

The model tested showed optimal fit indexes: chi-squared ( $df = 4$ ) = 5.75,  $p = .218$ ; CFI = .99, TLI = .96, RMSEA = .052; SRMR = .042 (Figure 2). The model showed that (a) higher *maternal anxiety* at delivery was significantly associated with higher *prenatal stress* and lower *prenatal support*, (b) higher scores in postnatal *parenting stress* were significantly linked to higher *maternal anxiety* at delivery, (c) higher *mother–infant bonding* was significantly associated with higher *prenatal support* and lower *maternal anxiety* at delivery, and (d) infants' *regulatory capacity* was significantly linked with less *parenting stress* and more *mother–infant bonding* at 3 months. In this model, a total mediation of *maternal anxiety* emerged for the relationship between *prenatal stress* and *parenting stress* at 3 months, whereas a partial mediation emerged for *maternal anxiety* on the relationship between *prenatal support* and *mother–infant bonding*. In addition, a total mediation emerged for both *parenting stress* and *mother–infant bonding* on the relationship between *maternal anxiety* and infants' *regulatory capacity* at 3 months.

## Discussion

To the best of our knowledge, this is the first longitudinal and prospective study that documents the short-term consequences of COVID-19 pandemic-related prenatal maternal stress on infants' temperament (i.e., regulatory capacity) at 3 months. The model highlighted that a complex pathway may indirectly link prenatal stress experienced by women during the COVID-19 emergency with infants' regulatory capacity. The effect appeared to be mediated by different dimensions of maternal psychosocial well-being (i.e., anxiety and social support) and of the caregiving environment (i.e., parenting stress and bonding). More specifically, the model suggested that (a) high levels of pandemic-related prenatal stress and concurrent low levels of

**Table 2.** Descriptive statistics for women's physical exposure to COVID-19

During pregnancy...	Response	<i>N</i>	%
I tested positive for SARS-CoV-2	Yes	0	0
	No	163	100.0
I had symptoms reminiscent of COVID-19	Yes	16	9.8
	No	147	90.2
I had contacts with relatives or friends who tested positive for SARS-CoV-2	Yes	12	7.4
	No	151	92.6
I live in a high-contagion or hotspot area for COVID-19 spread (e.g., red zones)	Yes	51	31.3
	No	112	68.7
I had contacts with relatives or friends who live in high-contagion or hotspot areas	Yes	16	9.8
	No	147	90.2
One of my relatives or friends was hospitalized in an intensive care unit with COVID-19	Yes	39	23.9
	No	124	76.1
One of my relatives or friends died with COVID-19	Yes	24	14.7
	No	139	85.3

perceived social support may increase postnatal maternal anxiety, (b) higher postnatal maternal anxiety may increase maternal parenting stress and reduce maternal bonding, and (c) infants' regulatory capacity may be negatively affected by greater parenting stress and reduced postnatal bonding at 3 months.


The present findings are consistent with previous literature suggesting that prenatal stress may increase maternal anxiety symptoms in the postpartum period (Field, 2018). In addition, we confirm that, even during the time of a pandemic, the availability of social support may result in a protective effect capable of reducing postnatal maternal anxiety, replicating previous findings in community samples (Razurel, Kaiser, Sellenet, & Epiney, 2013). For pregnant women, the COVID-19 healthcare emergency may have resulted in increased prenatal stress due to worries related to contagion and reduced social support as a consequence of lockdown and social isolation. As such, it is not surprising that the mothers included in this sample presented high levels of anxiety, with about one out of four women reporting anxiety levels well above the cut-off for clinical severity. It is noteworthy that there were no statistically significant differences in prenatal stress and postpartum anxiety between mothers with and without direct or indirect exposures to COVID-19. In other words, developing COVID-19 symptomatology and the close relationship with significant ones who were positive for COVID-19, required intensive care hospitalization or died with the disease were not factors associated with significant increases in psychological symptoms. This finding suggests that protecting and supporting maternal psychological health during and after the pandemic should be a priority goal for clinicians and policy makers. Moreover, the risk of heightened anxiety should not be underestimated, even in low-risk community samples.

The described increase in anxiety symptomatology emerged as a risk factor for heightened parenting stress and reduced maternal bonding 3 months after delivery. On the one hand, this finding is consistent with previous reports that have suggested how maternal

**Table 3.** Preliminary bivariate Pearson's correlations

	1	2	3	4	5	6
1. Physical exposure to COVID-19	1					
2. Prenatal COVID-19-related maternal stress	.032	1				
3. Prenatal maternal social support	.032	.040	1			
4. Postnatal maternal state anxiety	-.112	.218**	-.210**	1		
5. Postnatal maternal parenting stress	-.091	.199*	-.157*	.483**	1	
6. Postnatal mother–infant bonding	.008	-.073	.258**	-.399**	-.602**	1
7. Infant regulatory capacity	.003	.002	.227**	-.189*	-.328**	.312**

Note: \* $p < .05$ ; \*\* $p < .01$ .

**Figure 2.** Path model of the relationships between the study variables.

postnatal anxiety may be significantly associated with specific affective and emotional dimensions of caregiving (Müller et al., 2016; Riva Crugnola et al., 2016). On the other hand, this further suggests that the exposure to the stress related to the present pandemic may indirectly result in a less-than-optimal caregiving environment characterized by high levels of parenting stress and specific challenges in developing a sense of intimate closeness and attachment to the infant. Of note, a direct protective effect of social support emerged on maternal bonding. Previous research suggests that interventions aimed at providing pregnant women and mothers with social support may increase maternal feelings of bonding and emotional closeness in the first months after delivery (Ohara et al., 2017). Thus, it seems obvious to suggest that preventive interventions during the present healthcare emergency should be aimed to improve the quality of the caregiving environment, facilitating and promoting a better sense of affective bonding between mothers and infants and reducing the risk of heightened parenting stress.

Finally, infants' temperament was indirectly affected by the heightened maternal stress experienced during the present pandemic as high levels of parenting stress and reduced maternal bonding appeared to be significantly associated with lower scores

in infants' regulatory capacity at 3 months. Infants' regulatory skills are highly susceptible to alterations in the caregiving environment (Davis & Narayan, 2020; Gartstein & Skinner, 2018). Prenatal stress is especially a risk factor for altered fetal development that may end up in a dysregulated temperament profile in infancy and childhood (Howland, Sandman, Davis, & Glynn, 2020). Consistently, infants born during the COVID-19 healthcare emergency may be at heightened risk for developing less than optimal regulatory skills and this seems to be indirectly affected by a cascade effect of pandemic-related stress on maternal stress, anxiety, and bonding. As the effect of prenatal stress on infants' regulatory capacities passed through the alterations in the caregiving environment, these findings also highlight the opportunity to invest in family-centered care strategies during and after the present pandemic to provide mothers and infant with optimal support. The prenatal stress to which pregnant women were exposed during the COVID-19 emergency may have set a less-than-optimal trajectory for infants' temperament development. At the same time, it is plausible to hypothesize that effective protective interventions may be successful as far as they target the same primary predictors of infants' regulatory capacity: maternal stress and bonding. By investing in relational

interventions, clinicians and policy makers may create opportunities for healing togetherness even in a time of crisis and separation.

### Limitations

The sample size was relatively small and did not allow the inclusion of multiple mediators and confounders in the model. We collected data using only self-report instruments and the maternal retrospective report about the prenatal stress experienced may be partially affected by a recall bias. The COVID-19-related prenatal stress questionnaire was developed ad hoc for this study and there is no standardization. However, this methodological choice allowed us to obtain a measure of pandemic-related stress, whereas a more general self-report questionnaire on prenatal stress would have been too broad and less specific to the contingency of the present healthcare emergency. In addition, to limit the exposure to pandemic-related stress and not to the COVID-19 disease itself, we opted to include only mothers who tested negative to SARS-CoV-2 at delivery. The test was done in all the neonatal units using nasopharyngeal polymerase chain reaction, but serology was not performed systematically. To further control for this, we included specific items in the COVID-19 exposure questionnaire that asked the mothers to report if they were positive for the SARS-CoV-2 during pregnancy and if they had symptoms reminiscent of COVID-19. No woman reported to have tested positive for SARS-CoV-2 during pregnancy. Through a conservative set of inclusion criteria, we excluded and controlled additional sources of stress; nonetheless, other sources should not be excluded. For example, we did not control for the number of previous children already present in the family or other traumatic experiences that may have occurred in women's childhood. Finally, all the enrolled mother–infant dyads lived in northern Italy and the findings need replications in other mother–infant populations.

### Conclusions and implications

The present study contributes to understanding how the COVID-19 healthcare emergency may be affecting infants' development through prenatal exposure to maternal pandemic-related stress, even in a low-risk community sample. The findings outline the risk of a hidden pandemic of developmental psychopathology that involves both maternal stress and anxiety, as well as the risk of infants' temperamental dysregulation. The findings have implications for scientific advances and policy-maker decisions relevant for maternal and pediatric healthcare.

The indirect pathway highlighted in this study leaves open the question related to the mechanisms involved in the transmission of prenatal maternal pandemic-related stress to infants' behavioral development. Recent research suggests that environmental exposures to stress that are capable of altering the caregiving environment may lead to detrimental effects for infants' behavioral development through epigenetic mechanisms. Increased DNA methylation is such an epigenetic marker that is highly susceptible to environmental adversities and that may affect infants' temperament, especially through effects on the transcriptional activity of stress-related genes (Gartstein & Skinner, 2018). Variations in the maternal psychosocial well-being and the caregiving environment have been linked with altered infants' methylation of glucocorticoid receptor gene (nuclear receptor subfamily 3, group C, member 1 [*NR3C1*]), serotonin transporter gene (solute carrier

family C6, member 4 [*SLC6A4*]), and other candidate loci involved in socioemotional and stress reactivity (Berretta, Guida, Forni, & Provenzi, 2021; Devlin, Brain, Austin, & Oberlander, 2010; Oberlander et al., 2008; Provenzi, Brambilla, Scotto di Minico, Montirosso, & Borgatti, 2020b). The MOM-COPE project also includes the assessment of DNA methylation in specific stress-related genes (e.g., *NR3C1* and *SLC6A4*). As such, future analyses from this cohort may further contribute to highlight the biological mechanisms that are involved in the complex indirect pathway that leads from pandemic-related prenatal stress to adverse effects on infants' behavioral development.

The present COVID-19 healthcare emergency may pose serious risks for infants' temperament development by setting the stage for a complex domino reaction triggered by increased pandemic-related stress and reduced social support. From this point of view, COVID-19-related prenatal stress appears to act as a hidden pandemic that should not be underestimated as it is affecting the early developmental trajectories of infants' behavioral development as early as the age of 3 months. The indirect effect of pandemic-related stress on infants' behavioral development may contribute to making the detrimental effects of maternal distress during the COVID-19 healthcare emergency less immediately visible and acknowledgeable by clinicians and by mothers themselves. Nonetheless, the present study signals the presence of a relevant risk for mother–infant health and highlights specific targets for potential interventions. Even in low-risk community families, the protection and promotion of mother–infant well-being should be prioritized during a time of pandemic (Provenzi, Baroffio, Ligabue, & Borgatti, 2020a). In this scenario, the present pandemic represents a unique opportunity to invest in relational preventive interventions from the very beginning of life, starting new programs for mother–child health or strengthening the existing ones. Such dedicated investment by policy makers is warranted to promote a culture of family-centered care that integrates relational and developmental psychopathology dimensions in routine follow-up assessments of infants' development across the first years of life. By taking care of early signals of regulatory issues in infants born during the COVID-19 emergency we may be able to prevent further behavioral problems in childhood, with benefits for both families and healthcare systems.

**Supplementary Material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0954579421000766>

**Acknowledgments.** Special thanks to Drs. Beril Calgan, Eleonora Fullone, Vanessa Manfredini, Francesca Masoni, Giada Pectenati, Elia Rinaldi, and Luisa Vercellino: they were trainees in psychology (IRCCS Mondino Foundation) at the time of study and they provided key support to data collection. Dr. Cinzia Fattore provided essential administrative support to the management of the multi-center collaborations involved in the MOM-COPE project during the difficult pandemic period and contributed to making this project possible. The authors are thankful to the families who participated in this study.

**Funding Statement.** This study is supported by funds from Roche Italy and from the Italian Ministry of Health (Cinque per Mille 2017) to author LP.

**Conflicts of Interest.** None.

### References

- Abidin, R. R. (1995). *Parenting stress index: Manual* (3rd ed.). Odessa, FL: Psychological Assessment Resources.
- Barello, S., Palamenghi, L., & Graffigna, G. (2020). Burnout and somatic symptoms among frontline healthcare professionals at the peak of the Italian

- COVID-19 pandemic. *Psychiatry Research*, 290, 113129. doi:10.1016/j.psychres.2020.113129
- Berretta, E., Guida, E., Forni, D., & Provenzi, L. (2021). Glucocorticoid receptor gene (NR3C1) methylation during the first thousand days: Environmental exposures and developmental outcomes. *Neuroscience and Biobehavioral Reviews*, 125, 493–502. doi:10.1016/j.neubiorev.2021.03.003
- Breslin, N., Baptiste, C., Gyamfi-Bannerman, C., Miller, R., Martinez, R., Bernstein, K., ... Goffman, D. (2020). Coronavirus disease 2019 infection among asymptomatic and symptomatic pregnant women: Two weeks of confirmed presentations to an affiliated pair of New York city hospitals. *American Journal of Obstetrics & Gynecology*, 2, 100118. doi:10.1016/j.ajogmf.2020.100118
- Brock, R. L., O'Hara, M. W., Hart, K. J., McCabe-Beane, J. E., Williamson, J. A., Brunet, A., ... King, S. (2015). Peritraumatic distress mediates the effect of severity of disaster exposure on perinatal depression: The Iowa flood study. *Journal of Traumatic Stress*, 28, 515–522. doi:10.1002/jts.22056
- Cameron, E. E., Joyce, K. M., Delaquis, C. P., Reynolds, K., Protudjer, J., & Roos, L. E. (2020). Maternal psychological distress & mental health service use during the COVID-19 pandemic. *Journal of Affective Disorders*, 276, 765–774. doi:10.1016/j.jad.2020.07.081
- Class, Q. A., Abel, K. M., Khashan, A. S., Rickert, M. E., Dalman, C., Larsson, H., ... D'Onofrio, B. M. (2014). Offspring psychopathology following pre-conception, prenatal and postnatal maternal bereavement stress. *Psychological Medicine*, 44, 71–84. doi:10.1017/S0033291713000780
- Condon, J. T., & Corkindale, C. J. (1998). The assessment of parent-to-infant attachment: Development of a self-report questionnaire instrument. *Journal of Reproductive and Infant Psychology*, 16, 57–76. doi:10.1080/02646839808404558
- Daglar, G., & Nur, N. (2018). Level of mother-baby bonding and influencing factors during pregnancy and postpartum period. *Psychiatria Danubina*, 30, 433–440. doi:10.24869/psyd.2018.433
- Davis, E. P., & Narayan, A. J. (2020). Pregnancy as a period of risk, adaptation, and resilience for mothers and infants. *Development and Psychopathology*, 32, 1625–1639. doi:10.1017/S0954579420001121
- Devlin, A. M., Brain, U., Austin, J., & Oberlander, T. F. (2010). Prenatal exposure to maternal depressed mood and the MTHFR c677 T variant affect SLC6A4 methylation in infants at birth. *PLoS One*, 5, e12201. doi:10.1371/journal.pone.0012201
- Eisenberg, N., Valiente, C., Spinrad, T. L., Liew, J., Zhou, Q., Losoya, S. H., ... Cumberland, A. (2009). Longitudinal relations of children's effortful control, impulsivity, and negative emotionality to their externalizing, internalizing, and co-occurring behavior problems. *Developmental Psychology*, 45, 988–1008. doi:10.1037/a0016213
- Fallon, V., Silverio, S. A., Halford, J., Bennett, K. M., & Harrold, J. A. (2021). Postpartum-specific anxiety and maternal bonding: Further evidence to support the use of childbearing specific mood tools. *Journal of Reproductive and Infant Psychology*, 39, 114–124. doi:10.1080/02646838.2019.1680960
- Feldman, R. (2015). Mutual influences between child emotion regulation and parent-child reciprocity support development across the first 10 years of life: Implications for developmental psychopathology. *Development and Psychopathology*, 27, 1007–1023. doi:10.1017/S0954579415000656
- Field, T. (2018). Postnatal anxiety prevalence, predictors and effects on development: A narrative review. *Infant Behavior & Development*, 51, 24–32. doi:10.1016/j.infbeh.2018.02.005
- Fuller, A., Messito, M. J., Mendelsohn, A. L., Oyeku, S. O., & Gross, R. S. (2018). Prenatal maternal hardships and infant regulatory capacity at 10 months old in low-income Hispanic mother-infant pairs. *Academic Pediatrics*, 18, 897–904. doi:10.1016/j.acap.2018.04.134
- Gartstein, M. A., Putnam, S. P., & Rothbart, M. K. (2012). Etiology of pre-school behavior problems: Contributions of temperament attributes in early childhood. *Infant Mental Health Journal*, 33, 197–211. doi:10.1002/imhj.21312
- Gartstein, M. A., & Rothbart, M. K. (2003). Studying infant temperament via the revised infant behavior questionnaire. *Infant Behavior and Development*, 26, 64–86. doi:10.1016/S0163-6383(02)00169-8
- Gartstein, M. A., & Skinner, M. K. (2018). Prenatal influences on temperament development: The role of environmental epigenetics. *Development and Psychopathology*, 30, 1269–1303. doi:10.1017/S0954579417001730
- Gunning, M., Halligan, S. L., & Murray, L. (2013). Contributions of maternal and infant factors to infant responding to the still face paradigm: A longitudinal study. *Infant Behavior and Development*, 36, 319–328. doi:10.1016/j.infbeh.2013.02.003
- Gutteling, B. M., De Weerth, C., Willemsen-Swinkels, S. H. N., Huizink, A. C., Mulder, E. J. H., Visser, G. H. A., & Buitelaar, J. K. (2005). The effects of prenatal stress on temperament and problem behavior of 27-month-old toddlers. *European Child and Adolescent Psychiatry*, 14, 41–51. doi:10.1007/s00787-005-0435-1
- Howland, M. A., Sandman, C. A., Davis, E. P., & Glynn, L. M. (2020). Prenatal maternal psychological distress and fetal developmental trajectories: Associations with infant temperament. *Development and Psychopathology*, 32, 1685–1695. doi:10.1017/S095457942000142X
- Huizink, A. C., Robles De Medina, P. G., Mulder, E. J. H., Visser, G. H. A., & Buitelaar, J. K. (2002). Psychological measures of prenatal stress as predictors of infant temperament. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41, 1078–1085. doi:10.1097/0004583-200209000-00008
- Laplante, D. P., Brunet, A., & King, S. (2016). The effects of maternal stress and illness during pregnancy on infant temperament: Project ice storm. *Pediatric Research*, 79, 107–113. doi:10.1038/pr.2015.177
- Lebel, C., MacKinnon, A., Bagshawe, M., Tomfohr-Madsen, L., & Giesbrecht, G. (2020). Elevated depression and anxiety symptoms among pregnant individuals during the COVID-19 pandemic. *Journal of Affective Disorders*, 277, 5–13. doi:10.1016/j.jad.2020.07.126
- Lin, B., Crnic, K. A., Luecken, L. J., & Gonzales, N. A. (2014). Maternal prenatal stress and infant regulatory capacity in Mexican Americans. *Infant Behavior and Development*, 37, 571–582. doi:10.1016/j.infbeh.2014.07.001
- Lista, G., & Bresesti, I. (2020). Fatherhood during the COVID-19 pandemic: An unexpected turnaround. *Early Human Development*, 144, 105048. doi:10.1016/j.earlhumdev.2020.105048
- López-Morales, H., del Valle, M. V., Canet-Juric, L., Andrés, M. L., Galli, J. I., Poó, F., & Urquijo, S. (2021). Mental health of pregnant women during the COVID-19 pandemic: A longitudinal study. *Psychiatry Research*, 295, 113567. doi:10.1016/j.psychres.2020.113567
- Masiero, M., Mazzocco, K., Harnois, C., Cropley, M., & Pravettoni, G. (2020). From individual to social trauma: Sources of everyday trauma in Italy, the US and UK during the Covid-19 pandemic. *Journal of Trauma and Dissociation*, 21, 1–7. doi:10.1080/15299732.2020.1787296
- Matvienko-Sikar, K., Murphy, G., & Murphy, M. (2018). The role of prenatal, obstetric, and post-partum factors in the parenting stress of mothers and fathers of 9-month old infants. *Journal of Psychosomatic Obstetrics and Gynecology*, 39, 47–55. doi:10.1080/0167482X.2017.1286641
- Morikawa, M., Okada, T., Ando, M., Aleksic, B., Kunimoto, S., Nakamura, Y., ... Ozaki, N. (2015). Relationship between social support during pregnancy and postpartum depressive state: A prospective cohort study. *Scientific Reports*, 5, 1–9. doi:10.1038/srep10520
- Müller, M., Tronick, E., Zietlow, A. L., Nonnenmacher, N., Verschoor, S., & Träuble, B. (2016). Effects of maternal anxiety disorders on infant self-comforting behaviors: The role of maternal bonding, infant gender and age. *Psychopathology*, 49, 295–304. doi:10.1159/000448404
- Nicol-Harper, R., Harvey, A. G., & Stein, A. (2007). Interactions between mothers and infants: Impact of maternal anxiety. *Infant Behavior and Development*, 30, 161–167. doi:10.1016/j.infbeh.2006.08.005
- Oberlander, T. F., Weinberg, J., Papsdorf, M., Grunau, R., Misri, S., & Devlin, A. M. (2008). Prenatal exposure to maternal depression, neonatal methylation of human glucocorticoid receptor gene (NR3C1) and infant cortisol stress responses. *Epigenetics*, 3, 97–106. doi:10.4161/epi.3.2.6034
- Obrochta, C. A., Chambers, C., & Bandoli, G. (2020). Psychological distress in pregnancy and postpartum. *Women and Birth*, 33, 583–591. doi:10.1016/j.wombi.2020.01.009
- Ohara, M., Okada, T., Aleksic, B., Morikawa, M., Kubota, C., Nakamura, Y., ... Ozaki, N. (2017). Social support helps protect against perinatal bonding failure and depression among mothers: A prospective cohort study. *Scientific Reports*, 7, 9546. doi:10.1038/s41598-017-08768-3



- Pope, J., Olander, E. K., Leitao, S., Meaney, S., & Matvienko-Sikar, K. (2021). Prenatal stress, health, and health behaviours during the COVID-19 pandemic: An international survey. *Women and Birth*. ahead of print. doi:10.1016/j.wombi.2021.03.007
- Posner, M. I., & Rothbart, M. K. (2000). Developing mechanisms of self-regulation. *Development and Psychopathology*, *12*, 427–441. doi:10.1017/S0954579400003096
- Preis, H., Mahaffey, B., Heiselman, C., & Lobel, M. (2020). Vulnerability and resilience to pandemic-related stress among U.S. Women pregnant at the start of the COVID-19 pandemic. *Social Science and Medicine*, *266*, 113348. doi:10.1016/j.socscimed.2020.113348
- Provenzi, L., Baroffio, E., Ligabue, S., & Borgatti, R. (2020a). The little professor and the virus: Scaffolding children's meaning making during the COVID-19 emergency. *Frontiers in psychiatry*, *11*, 817. doi:10.3389/fpsy.2020.00817
- Provenzi, L., Brambilla, M., Scotto di Minico, G., Montiroso, R., & Borgatti, R. (2020b). Maternal caregiving and DNA methylation in human infants and children: Systematic review. *Genes, Brain and Behavior*, *19*, e12616. doi:10.1111/gbb.12616
- Provenzi, L., Fumagalli, M., Bernasconi, F., Sirgiovanni, I., Morandi, F., Borgatti, R., & Montiroso, R. (2017). Very preterm and full-term infants' response to socio-emotional stress: The role of postnatal maternal bonding. *Infancy*, *22*, 695–712. doi:10.1111/infa.12175
- Provenzi, L., Grumi, S., Giorda, R., Biasucci, G., Bonini, R., Cavallini, A., ... Borgatti, R. (2020c). Measuring the outcomes of maternal COVID-19-related prenatal exposure (MOM-COPE): Study protocol for a multicentric longitudinal project. *BMJ Open*, *10*, doi:10.1136/bmjopen-2020-044585
- Provenzi, L., & Tronick, E. (2020). The power of disconnection during the COVID-19 emergency: From isolation to reparation. *Psychological Trauma: Theory, Research, Practice, and Policy*, *12*, S252–S254. doi:10.1037/tra0000619
- Racine, N., Devereaux, C., Cooke, J. E., Eirich, R., Zhu, J., & Madigan, S. (2021). Adverse childhood experiences and maternal anxiety and depression: A meta-analysis. *BMC Psychiatry*, *21*, 28. doi:10.1186/s12888-020-03017-w
- Racine, N., Plamondon, A., Hentges, R., Tough, S., & Madigan, S. (2019). Dynamic and bidirectional associations between maternal stress, anxiety, and social support: The critical role of partner and family support. *Journal of Affective Disorders*, *252*, 19–24. doi:10.1016/j.jad.2019.03.083
- Razurel, C., Kaiser, B., Sellenet, C., & Epiney, M. (2013). Relation between perceived stress, social support, and coping strategies and maternal well-being: A review of the literature. *Women and Health*, *53*, 74–99. doi:10.1080/03630242.2012.732681
- R Core Team. (2020). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>.
- Riva Crugnola, C., Ierardi, E., Ferro, V., Gallucci, M., Parodi, C., & Astengo, M. (2016). Mother-infant emotion regulation at three months: The role of maternal anxiety, depression and parenting stress. *Psychopathology*, *49*, 285–294. doi:10.1159/000446811
- Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, *48*, 1–36. doi:10.18637/jss.v048.i02
- Salehi, L., Rahimzadeh, M., Molaei, E., Zaheri, H., & Esmaelzadeh-Saeieh, S. (2020). The relationship among fear and anxiety of COVID-19, pregnancy experience, and mental health disorder in pregnant women: A structural equation model. *Brain and Behavior*, *10*, e01835. doi:10.1002/brb3.1835
- Schwartz, D. A. (2020). An analysis of 38 pregnant women with COVID-19, their newborn infants, and maternal-fetal transmission of SARS-CoV-2: Maternal coronavirus infections and pregnancy outcomes. *Archives of Pathology and Laboratory Medicine*, *144*, 799–805. doi:10.5858/arpa.2020-0901-SA
- Sheinkopf, S. J., Lester, B. M., LaGasse, L. L., Seifer, R., Bauer, C. R., Shankaran, S., ... Wright, L. L. (2006). Interactions between maternal characteristics and neonatal behavior in the prediction of parenting stress and perception of infant temperament. *Journal of Pediatric Psychology*, *31*, 27–40. doi:10.1093/jpepsy/jsj026
- Simcock, G., Laplante, D. P., Elgbeili, G., Kildea, S., Cobham, V., Stapleton, H., & King, S. (2017). Infant neurodevelopment is affected by prenatal maternal stress: The QF2011 Queensland flood study. *Infancy*, *22*, 282–302. doi:10.1111/infa.12166
- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. (1983). *Manual for the state-trait anxiety inventory*. Palo Alto, CA: Consulting Psychologists Press.
- Su, Q., Zhang, H., Zhang, Y., Zhang, H., Ding, D., Zeng, J., ... Li, H. (2015). Maternal stress in gestation: Birth outcomes and stress-related hormone response of the neonates. *Pediatrics and Neonatology*, *56*, 376–381. doi:10.1016/j.pedneo.2015.02.002
- Tani, F., & Castagna, V. (2017). Maternal social support, quality of birth experience, and post-partum depression in primiparous women. *Journal of Maternal-Fetal and Neonatal Medicine*, *30*, 689–692. doi:10.1080/14767058.2016.1182980
- Tietz, A., Zietlow, A. L., & Reck, C. (2014). Maternal bonding in mothers with postpartum anxiety disorder: The crucial role of subclinical depressive symptoms and maternal avoidance behaviour. *Archives of Women's Mental Health*, *17*, 433–442. doi:10.1007/s00737-014-0423-x
- Van Bussel, J. C. H., Spitz, B., & Demyttenaere, K. (2010). Three self-report questionnaires of the early mother-to-infant bond: Reliability and validity of the Dutch version of the MPAS. *PBQ and MIBS*. *Archives of Women's Mental Health*, *13*, 373–384. doi:10.1007/s00737-009-0140-z
- Van den Bergh, B. R. H., van den Heuvel, M. I., Lahti, M., Braeken, M., de Rooij, S. R., Entringer, S., ... Schwab, M. (2020). Prenatal developmental origins of behavior and mental health: The influence of maternal stress in pregnancy. *Neuroscience and Biobehavioral Reviews*, *117*, 26–64. doi:10.1016/j.neubiorev.2017.07.003
- Wittig, S. M. O., & Rodriguez, C. M. (2019). Interaction between maternal and paternal parenting styles with infant temperament in emerging behavior problems. *Infant Behavior and Development*, *57*, 101323. doi:10.1016/j.infbeh.2019.04.005
- Xie, R. H., He, G., Koszycki, D., Walker, M., & Wen, S. W. (2009). Prenatal social support, postnatal social support, and postpartum depression. *Annals of Epidemiology*, *19*, 637–643. doi:10.1016/j.annepidem.2009.03.008
- Xie, R.-H., Yang, J., Liao, S., Xie, H., Walker, M., & Wen, S. W. (2010). Prenatal family support, postnatal family support and postpartum depression. *Australian and New Zealand Journal of Obstetrics and Gynaecology*, *50*, 340–345. doi:10.1111/j.1479-828X.2010.01185.x
- Xiong, J., Lipsitz, O., Nasri, F., Lui, L. M. W., Gill, H., Phan, L., ... McIntyre, R. S. (2020). Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *Journal of Affective Disorders*, *277*, 55–64. doi:10.1016/j.jad.2020.08.001
- Zaigham, M., & Andersson, O. (2020). Maternal and perinatal outcomes with COVID-19: A systematic review of 108 pregnancies. *Acta Obstetrica et Gynecologica Scandinavica*, *99*, 823–829. doi:10.1111/aogs.13867
- Zhu, P., Sun, M.-S., Hao, J.-H., Chen, Y.-J., Jiang, X.-M., Tao, R.-X., ... Tao, F.-B. (2014). Does prenatal maternal stress impair cognitive development and alter temperament characteristics in toddlers with healthy birth outcomes? *Developmental Medicine & Child Neurology*, *56*, 283–289. doi:10.1111/dmcn.12378
- Zimet, G. D., Dahlem, N. W., Zimet, S. G., & Farley, G. K. (1988). The multidimensional scale of perceived social support. *Journal of Personality Assessment*, *52*, 30–41. doi:10.1207/s15327752jpa5201\_2