

Short Report

Cite this article: Cena L, Trainini A, Belluardo M, Buizza C, Cooper H, Murray L. (2026) Dialogic Book-Sharing intervention in an Italian multicentre randomized controlled trial: effects on parental mood. *Primary Health Care Research & Development* 27(e82): 1–4. doi: [10.1017/S1463423626101509](https://doi.org/10.1017/S1463423626101509)

Received: 4 September 2025 UTC

Revised: 5 June 2026 UTC

Accepted: 25 June 2026 UTC

Keywords:

anxiety symptoms; depression symptoms; Dialogic Book-Sharing; parent-child early intervention; parental mood

Corresponding author:

Loredana Cena; Email: loredana.cena@unibs.it

Alice Trainini and Mauro Belluardo are contributed equally to this work.

© The Author(s), 2026. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press or the rights holder(s) must be obtained prior to any commercial use and/or adaptation of the article.



Dialogic Book-Sharing intervention in an Italian multicentre randomized controlled trial: effects on parental mood

Loredana Cena¹ , Alice Trainini¹ , Mauro Belluardo², Chiara Buizza¹ , Hal Cooper³ and Lynne Murray⁴

¹Observatory of Perinatal Clinical Psychology, Department of Clinical and Experimental Sciences, Brescia University, Italy; ²Department of Medicine and Surgery, University of Parma: Università degli Studi di Parma, Italy; ³Oxford University: University of Oxford, UK and ⁴School of Psychology and Clinical Language Sciences, University of Reading, UK

Abstract

This randomized controlled trial (RCT) was conducted in 10 Italian health and maternal-child Centres to evaluate the effects of a Dialogic Book-Sharing (DBS) intervention on parental symptoms of anxiety and depression. Sixty-eight parents of 14–20-month-old infants were recruited. ANOVA analysis revealed a greater reduction in anxiety and depression symptoms in the intervention group ($N = 36$) compared to controls ($N = 32$). DBS is a simple, cost-effective programme that benefits parental mood, making it a valuable intervention for perinatal populations.

Introduction

Clinical Psychology has highlighted how early experiences are important for child development and how these occur mainly in the context of parent-child relationships. Negative parental affective states, particularly depression and anxiety, have an adverse impact on parent-child interactions and child development. Thus, offspring of postnatally depressed mothers evidence child attachment insecurity, negative affective outcomes in childhood (assessed at 8 years) and adolescence (assessed at 13 and 16 years), and compromised cognitive and behavioural development (Murray *et al.*, 2011). Infants of anxious mothers show heightened stress responses, lower social engagement, and are twice as likely to develop anxiety as children of non-anxious mothers (Lawrence *et al.*, 2019).

Early parenting interventions have been shown to improve child outcomes. For example, parent-child play and shared reading interventions enhance children's cognitive and linguistic skills as well as their social-emotional development, and reduce externalizing behaviours at 36 months (Weisleder *et al.*, 2019). Far less is known, however, about the impact of such programmes on parents, and particularly their mental state. Although a meta-analysis of shared reading programmes involving parents and children aged 0–6 years old concluded that they may have some positive impact on parents' psychosocial functioning, only three of the included studies assessed parental depression, and none identified a benefit (Xie *et al.*, 2018).

A structured parent-child shared reading intervention principally directed at improving children's cognitive development in early childhood was developed by Murray and Cooper (Cooper *et al.*, 2014), the 'Dialogic Book-Sharing' (DBS) programme. This intervention has been evaluated in several countries including South Africa with children aged 23 and 27 months (Dowdall *et al.*, 2017), Lesotho with 1–5 years old children (Tomlinson *et al.*, 2016), Brazil with children aged between 30 and 42 months (Murray *et al.*, 2019), and England with children between 28 and 45 months of age (Murray *et al.*, 2018) with positive effects on both parenting and children's development. A consistent finding across all the studies is significant benefits for caregiver-child interactions during book-sharing, particularly in terms of parental *sensitivity* (appropriate and warm responsiveness to the child) and parent-child *reciprocity*.

DBS creates a special 'intersubjective' space (Murray *et al.*, 2022) where both parent and child share visual focus in close proximity, fostering shared physiological and emotionally positive states (Waters *et al.*, 2017). The emotional cues transmitted by the caregiver during these interactions help imprint the relational experience in memory, and can play a role in fostering secure attachment between parent and child (Bus and Van IJzendoorn, 1997). Despite evidence for benefits to caregiver-child interactions, there is limited research on whether this positive impact extends to caregivers more broadly, including their mental state. However, notwithstanding the aforementioned meta-analytic evidence that did not demonstrate benefit for parental depression (Xie *et al.*, 2018), more recent evidence points to potential for shared reading to have positive effects on parent outcomes. For example, in a mediation analysis examining a pathway from shared reading to child outcomes via parent psychosocial



functioning, Weisleder *et al.* (2019) found significant effects of a DBS intervention on parent stress about the parent-child relationship, and on maternal depressive symptoms. Further, in a longitudinal secondary analysis of data from an RCT, Canfield *et al.* (2020) found that levels of observationally assessed DBS quality at 6-months were associated with higher relational health (indicated by parental warmth and sensitivity) and lower parenting stress at 18-months. Taken together, there is a conceivable mechanism by which DBS may improve parent-child relations and parenting stress, and that this may in turn positively impact parent mental health outcomes.

Given the limited evidence for DBS's effects on parent mental health outcomes to date, the Observatory of Perinatal Clinical Psychology of the University of Brescia examined the application of the DBS in Italy using a randomized controlled trial (RCT) to assess the effects of this intervention on infant development, the quality of the parent-child relationship, and parental mental state. In the current paper we report on the impact of a DBS programme delivered to parents on parental symptoms of anxiety and depression.

Materials and methods

Study design

This RCT is part of a larger project in Italy (Cena *et al.*, 2024) promoted by the Observatory of Perinatal Clinical Psychology. Health, educational and maternal-child centres in Italy took part in a multicentre, two parallel arms, single blinded randomized controlled superiority trial, with a 1:1 arms allocation ratio on the impact of a DBS training provided to parents. Information from a centre in each of 10 cities (Brescia, Bergamo, Milan, Mantua, Como, La Spezia, Florence, Rome, Naples, Salerno) comprise the data set.

The intervention

The core feature of the DBS programme is showing parents how to support their child's interest and active engagement, rather than simply 'reading' to their child. Responsiveness is emphasized that is sensitive to the child's developmental capacity and experience, as well as the importance of a positive encouraging approach. The intervention was delivered by local staff who had been trained to act as facilitators over a two-day course (run by MB). Facilitators received ongoing supervision from LM in remote sessions that ensured fidelity with programme delivery. The intervention programme was delivered within each Centre in Italian, using Powerpoint slides and embedded demonstration videos. The programme was structured into four, weekly, meetings of the parents lasting 90 minutes each, which took place at the maternal-child/educational centres. The sessions were conducted with groups of up to six parents (mothers and fathers), during which children were cared for in an adjoining play space by Centre staff. Group sessions were followed by a brief period of one-to-one discussion between the facilitator and each parent (together with their child). Four illustrated wordless books were used. Parents were encouraged to apply the session's techniques during the following week in their daily life with their children.

Participants and ethics

Recruitment took place on a voluntary basis in the aforementioned centres by local staff. Inclusion criteria were: being able to speak

and read Italian; and having a child aged 14–20 months. The exclusion criteria were: parent having a diagnosed medical condition that could compromise their ability to participate in the intervention programme; or child having a diagnosed physical or intellectual impairment.

At each centre, parents were randomly assigned to either the intervention or the control condition using a random blocks algorithm. The intervention group received the DBS training immediately, while the control group received no input from the research team, apart from the assessment (controls received the DBS training after the final study assessment).

Data were collected one week before the intervention for baseline assessments and one week after the intervention period for follow up assessments. Local staff, not involved in the DBS delivery, blinded to group allocation, collected data before randomization and again after DBS delivery.

A total of 200 parents were invited by centre staff to take part in the study. Sixty-eight (34%) accepted. The reasons for non-acceptance were mainly lack of time, work commitments, and difficulties in family organization.

An Information Sheet with a description of the aims of the study was given to participants. To participate, the parent was asked to sign an Informed Consent Form. The protection of their privacy was ensured by the assignment of a code to the participants. The research followed the APA Ethical Principles of Psychologists and Code of Conduct.

The study was approved by the Institutional Ethics Board of the ASST Spedali Civili of Brescia, Italy (reference number NP5205 27.09.2022). All procedures performed in this study are in accordance with the ethical standards of the Institutional Ethics Board of the ASST Spedali Civili of Brescia, Italy and with the 1964 Declaration of Helsinki and its later amendments.

Instruments

A self-report questionnaire for parents taking approximately 15 minutes to complete was used for the assessment. It included:

- *Socio-demographic information Form*: sociodemographic data, and information about pregnancy and the post-partum period.
- *HADS Hospital Anxiety and Depression Scale* (Zigmond and Snaith, 1983): a 14-item self-report questionnaire assessing symptoms of anxiety and depression. Cronbach's alpha was 0.7 for the anxiety scale and 0.6 for depression scale.

Statistical analysis

Independent *t* tests and Chi square tests (or Fisher exact tests) were used to compare groups at baseline. Repeated measures ANOVAs were used to assess differences between the intervention and control groups over time; subsequently, non-independent post hoc *t* tests were used to assess change within each of the two groups across the two assessment points (the non-parametric Wilcoxon test was applied when normality assumptions were not met). All tests were two-tailed with $p < .05$.

Results

Sixty-eight parent-child dyads (36 experimental group and 32 control group) were recruited in the study and completed assessments at both baseline and follow up. No statistically significant differences were found between the two groups at baseline:

Table 1. Comparison between Index and Control groups scores

	Baseline					
	Intervention	Control	Statistics	p Value		
	N = 36	N = 32				
Parent's age (years), mean (SD)	34.2 (4.6)	36.4 (5.3)	1.564*	.90		
Parent's gender (mother)	32 (89.0%)	30 (93.7%)	.498 [#]	.48		
Italian nationality	32 (88.9%)	25 (78.1%)	11.665 [#]	.11		
University degree	13 (36.1%)	18 (56.3%)	8.724 [#]	.06		
Employed	25 (69.4%)	24 (75%)	5.303 [#]	.15		
High SES	15 (41.7%)	17 (53.1%)	4.405 [#]	.22		
Married/cohabiting	29 (80.6%)	25 (78.1%)	1.767 [#]	.62		
	Baseline		Post-intervention (after 4 weeks)		ANOVA	
	Intervention	Control	Intervention	Control	F ₂	p Value
	N = 36	N = 32	N = 36	N = 32		
Anxiety	4.9 (2.8)	4.9 (2.7)	4.1 (2.5)	5.6 (3.1)	5.327	.02
Depression	4.6 (3.0)	4.8 (3.1)	3.8 (2.8)	5.1 (3.2)	3.870	.05

*T test; [#]Chisquare.

they were similar with regard to demographic features, and were generally at low socio-economic risk (see Table 1). Baseline and post-intervention scores for the parental questionnaires are also shown in Table 1, together with statistical comparisons.

For parental anxiety symptoms, the ANOVA revealed a significant group/time interaction effect ($F_2 = 5.327$, $p = .02$, $\eta^2 = .09$). There was a similar finding of an interaction effect for parental depressive symptoms which fell just short of conventional statistical significance ($p = .05$, $\eta^2 = .06$). Post hoc within group analyses showed a significant decline in the intervention group for both anxiety ($p = .018$; $r = 0.5$) and depression ($p = .033$; $r = 0.4$); the change in scores in the control group, by contrast, was not significant for either measure (anxiety $p = .816$ and depression $p = .940$).

Discussion

Intervention and control groups differed from each other over time in their symptoms of anxiety and depression. For parents in the DBS intervention group, levels of both anxiety and depression showed a significant decrease from the week prior to the intervention to the week after the intervention was completed. The symptoms of parents in the control group, by contrast, showed no significant change over the same period.

The hypothesized mechanisms for this effect is that positive experiences improve parent-child relations and reduce stress, which is mood elevating for the parent.

This study had some limitations that should be acknowledged. First, the sample was composed predominantly of mothers, with relatively limited participation of fathers. As such, the findings may not fully capture potential differences in how parenting interventions affect maternal versus paternal mental health. Second, participants were generally at low socio-economic risk: this may limit the generalizability of the findings.

Although our study was small and therefore requires replication, our finding of a benefit to parents of DBS with their children is encouraging and adds to previous research showing that the DBS intervention carries benefits to parent-child interactions

and child development. DBS in early childhood is simple, practical, and inexpensive, making it suitable for widespread implementation in early childhood social and educational services. It can readily be integrated into educational programmes, providing valuable support for both parents and children. In sum, this evidence suggests that DBS is a suitable programme to not only promote child development, but also parental well-being. Future research on the effects of DBS should measure parent mental health outcomes, and should facilitate mediation analysis to examine the hypothesized pathway to promoting parent mental health via the reduction of parenting stress and the improvement of parent-child relationships.

Data availability statement. The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgements. We would like to thank the facilitators who were trained in the DBS intervention and the Directors who authorized its application in their health, educational and maternal-child centres in the Italian territory. We are also extremely grateful to all the parents and children who took part in this study.

Author contributions. Conceptualization, Loredana Cena and Lynne Murray; Data curation, Alice Trainini; Formal analysis, Chiara Buizza; Software, Chiara Buizza; Supervision, Lynne Murray and Mauro Belluardo; Writing – original draft, Loredana Cena and Alice Trainini; Writing – review & editing, Hal Cooper and Lynne Murray. All authors have read and agreed to the published version of the manuscript.

Funding statement. This work was funded by the Observatory of Perinatal Clinical Psychology (<https://www.unibs.it/it/node/988>) of the Department of Clinical and Experimental Science, University of Brescia (Italy).

Competing interests. The authors declare no conflicts of interest.

Institutional review board statement. The Study was reviewed and approved by the Institutional Ethics Board of the ASST Spedali Civili of Brescia, Italy (reference number NP5205 27.09.2022). All procedures performed in this study are in accordance with the ethical standards of the Institutional Ethics Board of the ASST Spedali Civili of Brescia, Italy and with the 1964 Declaration of Helsinki and its later amendments.

Informed consent statement. Before taking part in the study, all parents receive oral and written information about the intervention content and the assessment procedure. Informed consent was obtained from all subjects involved in the study.

References

- Bus AG and Van IJzendoorn MH (1997) Affective dimensions of mother-infant picture book reading. *Journal of School Psychology* 35(1), 47–60.
- Canfield CF, Miller EB, Shaw DS, Morris P, Alonso A and Mendelsohn AL (2020) Beyond language: Impacts of shared reading on parenting stress and early parent-child relational health. *Developmental Psychology* 56(7), 1305–1315.
- Cena L, Trainini A, Murray L, Cooper P, Calza S and Belluardo M (2024) Early parent-child intervention with Dialogic Book-Sharing: Effects on child communicative and socio-emotional development and on parenting. Study protocol for a multicentre randomised controlled trial in Italy. *Trials* 25(1), 395.
- Cooper PJ, Vally Z, Cooper H, Radford T, Sharples A, Tomlinson M and Murray L (2014) Promoting mother-infant book sharing and infant attention and language development in an impoverished South African population: A pilot study. *Early Childhood Education Journal* 42(2), 143–152.
- Dowdall N, Cooper PJ, Tomlinson M, Skeen S, Gardner F and Murray L (2017) The Benefits of Early Book Sharing (BEBS) for child cognitive and socio-emotional development in South Africa: Study protocol for a randomised controlled trial. *Trials* 18, 118.
- Lawrence PJ, Murayama K and Creswell C (2019) Systematic review and meta-analysis: Anxiety and depressive disorders in offspring of parents with anxiety disorders. *Journal of the American Academy of Child & Adolescent Psychiatry* 58(1), 46–60.
- Murray J, Santos IS, Bertoldi AD, Murray L, Arteché A, Tovo-Rodrigues L, Cruz S, Anselmi L, Martins R, Altafim E, Soares TB, Andriotti MG, Gonzalez A, Oliveira I, da Silveira MF and Cooper PJ (2019) The effects of two early parenting interventions on child aggression and risk for violence in Brazil (The PIÁ Trial): Protocol for a randomised controlled trial. *Trials* 20(1), 253.
- Murray L, Arteché A, Fearon P, Halligan S, Goodyer I and Cooper PJ (2011) Maternal postnatal depression and the development of depression in offspring up to 16 years of age. *Journal of the American Academy of Child & Adolescent Psychiatry* 50(5), 460–470.
- Murray L, Jennings S, Mortimer A, Prout A, Melhuish E, Hughes C, Duncan J, Holmes J, Dishington C and Cooper PJ (2018) The impact of early-years provision in Children Centres (EPICC) on child cognitive and socio-emotional development: Study protocol for a randomised controlled trial. *Trials* 19(1), 450.
- Murray L, Rayson H, Ferrari PF, Wass SV and Cooper PJ (2022) Dialogic Book-Sharing as a privileged intersubjective space. *Frontiers in Psychology* 13, 786991.
- Tomlinson M, Skeen S, Marlow M, Cluver L, Cooper P, Murray L, Mofokeng S, Morley N, Makhetha M, Gordon S, Esterhuizen T and Sherr L (2016) Improving early childhood care and development, HIV testing, treatment and support, and nutrition in Mokhotlong, Lesotho: A cluster randomized control trial of an integrated intervention. *Trials* 17, 538.
- Waters SF, West TV, Karnilowicz HR and Mendes WB (2017) Affect contagion between mothers and infants: Examining valence and touch. *Journal of Experimental Psychology: General* 146(7), 1043–1051.
- Weisleder A, Cates CB, Harding JF, Johnson SB, Canfield CF, Seery AM, Raak CD, Alonso A, Dreyer BP and Mendelsohn AL (2019) Links between shared reading and play, parent psychosocial functioning, and child behavior: Evidence from a randomized controlled trial. *The Journal of Pediatrics* 213, 187–195.e1.
- Xie QW, Chan CHY, Ji Q and Chan CLW (2018) Psychosocial effects of parent-child book reading interventions: A meta-analysis. *Pediatrics* 141, e20172675.
- Zigmond A and Snaith RP (1983) The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica* 67(6), 361–370.