

Arthroscopic treatment of an unusual distal clavicle osteochondroma causing rotator cuff impingement: case report and literature review

Piermarco Messinese, Valeria Vismara, Giuseppe Sircana, Vincenzo Campana, Fabrizio Mocini, Valentina Cardona, Andrea S. Monteleone, Rocco De Vitis, Giulio Maccauro, Maristella F. Saccomanno

Orthopaedic Institute, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

Abstract

Chronic shoulder impingement is one of the most common causes of shoulder pain. Intrinsic, extrinsic and secondary factors play a role in this syndrome; however the etiology of the pathology is still under debate. In rare cases, it can be caused by tumors, such as an osteochondroma. In the present study, a 49-year-old patient presented with shoulder pain for 6 months. Initially he underwent conservative treatment, without relief of symptoms. X-rays and MRI were then performed and showed the presence of an exostotic formation on the undersurface of the lateral third of the clavicle. The formation was arthroscopically removed. Histologic examination confirmed the diagnosis of osteochondroma. After surgery, the patient resumed fully activities with no symptoms within 3 months. At 1 year follow up, there are still no clinical or radiological signs of recurrence. This is, to our knowledge, the first case where an arthroscopic approach was used to remove an osteochondroma of the distal third of the clavicle.

Introduction

Chronic shoulder impingement is one of the most common causes of shoulder pain. It was first described by Neer, in 1972, as a pathological reduction of the subacromial space, which can cause a degeneration of the structures contained within.¹ It accounts for 44% to 65% of all shoulder girdle disorders,^{2,3} mainly occurring in overhead athletes and manual workers.⁴ The etiology of this syndrome is still under debate, since various different factors may play a role. Ellman *et al.* classified these factors in intrinsic, extrinsic, and secondary.⁵ Among

intrinsic factors, the natural process of aging of rotator cuff tendons,⁶ hypoxia, due to poor vascularity,⁷ and damage with tensile or shear loads,^{8,9} associated with overhead work and microtrauma, were found. On the other hand, extrinsic factors were linked to any morphological alterations, that causes a reduction in the subacromial space, and therefore compresses the supraspinatus tendon. Most common causes include: acromion morphology (especially Type III hooked described by Bigliani),¹⁰ acromial angle and slope,¹¹ acromial tilt, anterior acromial spur (mainly due to coracoclavicular ligament ossification), and acromioclavicular joint osteophytes.¹ Other morphological alterations that may cause this mechanical conflict are prominence of the greater tuberosity, thickening of subacromial bursa or the presence of an os acromialis. Biomechanical factors, instead, underlie secondary impingement mechanisms: scapula-humeral dyskinesia, muscle performance deficits, pectoralis minor and posterior capsule stiffness, postural abnormalities.^{12,13}

The aim of the present study was to describe a rare case of subacromial impingement, caused by an exostotic formation on the undersurface of the lateral third of the clavicle, which was treated arthroscopically.

Case Report

F.S. was a 49-year-old male with a progressive worsening of right shoulder pain and limitation of range of motion for the last 6 months. The patient did not have any previous known pathologies or traumas related to the shoulder, nor was the patient involved in overhead activities for work or sport. He presented to our clinic in April 2019 with a diagnosis of subacromial impingement, asking for a second opinion. He already underwent multiple sessions of physical therapy for the last 3 months, together with a medical therapy based of NSAID drugs, with no benefit.

At the clinical evaluation, no atrophy, deformity or sign of scapula dyskinesia could be detected. Passive range of motion was complete, although he reported pain upon elevation of the arm at 90°. Active range of motion was limited by pain. Impingement tests (Neer and Hawkins) were positive. No strength deficit could be noticed. X-rays showed an exostotic formation on the undersurface of the lateral third of the clavicle (Figure 1). Magnetic resonance (MR) clarified that the osseous formation abraded supraspinatus muscle belly. A layer of fluid between the osseous projection and the rotator

Correspondence: Piermarco Messinese, Via Ezio Sciamanna 113, 00168 Roma (RM), Italy.

Tel.: +39 327 7784636.

E-mail: piermarcomessinese@libero.it

Key words: subacromial impingement, osteochondroma, clavicle, arthroscopic surgery, rotator cuff.

Contributions: the authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none

Availability of data and materials: All data are reported in the text.

Ethics approval and consent to participate: Not applicable.

Informed consent: Informed consent has been obtained.

Received for publication: 11 April 2020.

Accepted for publication: 17 June 2020.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright: the Author(s), 2020

Licensee PAGEPress, Italy

Orthopedic Reviews 2020; 12(s1):8683

doi:10.4081/or.2020.8683

cuff was visualized. A thin rim of cartilage outlined the undersurface of the projection. The rotator cuff tendons were intact (Figure 2). Indication to an excisional biopsy was made. The procedure was performed arthroscopically. During surgery, no intra-articular pathology was found. In the subacromial space an abundant inflammatory bursitis was noticed. Rotator cuff tendons were intact. After bursectomy, the exostotic formation was found, consistently with the imaging, on the undersurface of the lateral third of the clavicle. It was arthroscopically resected to a level flush with the undersurface of the distal clavicle. After its removal, samples were sent for histology. Histologic examination confirmed the diagnosis of osteochondroma. After surgery, an immobilization sling was used for 7 days. Immediate passive range of motion was allowed. The patient resumed full activity, including sport, with no restriction or pain within 3 months after surgery.

Repeated postoperative imaging studies have revealed no recurrence at 6 and 12 months.

Discussion

Osteochondromas (OCs) account for 20-50% of all benign bony tumors, making it the most common among this class.¹⁴ In most cases, the OC arises during childhood, with a male predominance, and 75% of patients are younger than 20 years of age. Both solitary and multiple OCs can be found, even though the former is much more common. Multiple OCs are typical of patients affected by Multiple Hereditary Exostosis (HME). The OC is a surface lesion, which can be either pedunculated or sessile, composed both of cortical and medullary bone. A pathognomonic feature is the presence of cortical and medullary continuity, together with a cartilaginous cap. It can develop from any bone undergoing an enchondral maturation, but it is more commonly found at the level of long bones, especially those around the knee. Also flat bones can be affected, mainly the ilium and scapula.¹⁵

Clavicular OCs are rare, making up 0.2-0.5% of all solitary OCs.¹⁶ Only 11 cases of distal clavicle osteochondroma have been reported in the literature.¹⁷⁻²⁷ An open approach was usually used for the excision of the lesion. Of these, Galanopoulos *et al.*,²² Peidro *et al.*,¹⁸ Craig *et al.*²⁰ and Fallon *et al.*²¹ coupled en bloc resection with a distal clavicle excision. Simonetti *et al.*²⁷ tried an arthroscopic procedure, but, intraoperatively, due to the size of the lesion, decided to opt for an open surgery instead.

With the exception of two cases described by Alman *et al.*,²⁶ where the patients, 6 and 7-year-old males, did not experience any pain, in the vast majority of studies, impingement symptoms and decreased ROM were the most likely findings, in line with a compression of the supraspinatus tendon. Fallon *et al.*²¹ highlighted not only compression, but also a full thickness rotator cuff tear.

A medial clavicular localization of OC was instead described by Al Qudah *et al.*,²⁸ and Mallano *et al.*,²⁹ nonetheless, both opted for an open procedure.

A painful symptomatology can be observed not only when the neoformation was found at the level of the clavicle, but also at the level of the scapula.³⁰⁻³⁶ In fact, OCs have been found in this location in 3% to 4.6% of all cases.³⁷⁻³⁹ The lesions arose from different areas, such as the acromion,^{31-33,35,36} the coracoid, the superior part of the glenoid,³⁵ the costal surface near the superomedial angle of the scapula.³⁹ While Lopez Martin *et al.*³⁰ and Hommadi *et al.*³¹ chose an open surgical treatment, Citlack *et al.*³² started with an arthroscopic approach,

which had to be converted in mini-open, due to the size of the lesion. On the contrary, Lu *et al.*³³ and Simon Thomas *et al.*³⁴ used an arthroscopic procedure, avoiding the morbidity of an open resection. Main advantages of an arthroscopic approach are mainly related to avoidance of deltoid resection, decreased risk of infections,³³ together with an early functional recovery and cosmetic advantage.³⁸ In literature, just a few cases of shoulder OCs have been described. Although in most cases, the removal of the lesion was performed following an open approach, in this case study, instead, the lesion was completely removed arthroscopically, obtaining successful clinical results with no sign of recurrence at 1 year postop. The patient will

be followed up in time to make sure to early diagnose an eventual recurrence of disease. Surgery in this case was mainly indicated due to failure of conservative treatments for impingement symptoms, rather than due to the risk of malignant transformation, which in this kind of tumor is low, of about 1% when the lesion is solitary.^{14,40}

Conclusions

In conclusion, taking into account the size of the lesion and surgeon's experience and preference, the arthroscopic approach can be considered safe, effective and a viable option compared to open surgery.



Figure 1. X-rays, right shoulder antero-posterior view. An exostotic formation can be noticed on the undersurface of the lateral third of the clavicle.




Figure 2. Magnetic Resonance, right shoulder, sagittal view. A layer of fluid between the osseous projection and the rotator cuff can be visualized. A thin rim of cartilage outlined the undersurface of the projection.

References

1. Neer CS. Anterior acromioplasty for the chronic impingement syndrome in the shoulder: a preliminary report. *J Bone Joint Surg Am* 1972;54:41-50.
2. Feleus A, Bierma-Zeinstra SMA, Miedema HS, et al. Management in non-traumatic arm, neck and shoulder complaints: differences between diagnostic groups. *Eur Spine J Off Publ Eur Spine Soc Eur Spinal Deform Soc Eur Sect Cerv Spine Res Soc* 2008;17:1218-29.
3. Harrison AK, Flatow EL. Subacromial impingement syndrome. *J Am Acad Orthop Surg* 2011;19:701-8.
4. Zanca GG, Saccol MF, Oliveira AB, Mattiello SM. Shoulder internal and external rotations torque steadiness in overhead athletes with and without impingement symptoms. *J Sci Med Sport* 2013;16:433-7.
5. Ellman H, Gartsman GM. *Arthroscopic Shoulder Surgery and Related Procedures*. Malvern: Lea & Febiger, 1993.
6. Tempelhof S, Rupp S, Seil R. Age-related prevalence of rotator cuff tears in asymptomatic shoulders. *J Shoulder Elbow Surg* 1999;8:296-9.
7. Rudzki JR, Adler RS, Warren RF, et al. Contrast-enhanced ultrasound characterization of the vascularity of the rotator cuff tendon: age- and activity-related changes in the intact asymptomatic rotator cuff. *J Shoulder Elbow Surg* 2008;17:96S-100S.
8. Reilly P, Amis AA, Wallace AL, Emery RJH. Mechanical factors in the initiation and propagation of tears of the rotator cuff. Quantification of strains of the supraspinatus tendon in vitro. *J Bone Joint Surg Br* 2003;85:594-9.
9. Lake SP, Miller KS, Elliott DM, Soslowky LJ. Effect of fiber distribution and realignment on the nonlinear and inhomogeneous mechanical properties of human supraspinatus tendon under longitudinal tensile loading. *J Orthop Res Off Publ Orthop Res Soc* 2009;27:1596-602.
10. Bigliani LU, Ticker JB, Flatow EL, et al. The relationship of acromial architecture to rotator cuff disease. *Clin Sports Med* 1991;10:823-38.
11. Vaz S, Soyer J, Pries P, Clarac JP. Subacromial impingement: influence of coracoacromial arch geometry on shoulder function. *Joint Bone Spine* 2000;67:305-9.
12. Hallström E, Kärrholm J. Shoulder kinematics in 25 patients with impingement and 12 controls. *Clin Orthop* 2006;448:22-7.
13. Keener JD, Wei AS, Kim HM, et al. Proximal humeral migration in shoulders with symptomatic and asymptomatic rotator cuff tears. *J Bone Joint Surg Am* 2009;91:1405-13.
14. Murphey MD, Choi JJ, Kransdorf MJ, et al. Imaging of osteochondroma: variants and complications with radiologic-pathologic correlation. *Radiogr Rev Publ Radiol Soc N Am Inc* 2000;20:1407-34.
15. Motamedi K, Seeger LL. Benign bone tumors. *Radiol Clin North Am* 2011;49:1115-34.
16. Smith J, Yuppa F, Watson RC. Primary tumors and tumor-like lesions of the clavicle. *Skeletal Radiol* 1988;17:235-46.
17. Kontakis GM, Karantanas AH, Pasku D, et al. Delayed diagnosis of a symptomatic osteochondroma of the distal clavicle. *Orthopedics* 2006;29:734-6.
18. Peidro L, Suso S, Alcantara E, Ramon R. Periosteal chondroma of the clavicle. *Skeletal Radiol* 1996;25:406-8.
19. Vlychou M, Gibbons CLMH, Rigopoulou A. Bizarre parosteal osteochondromatous proliferation of the clavicle. *J Shoulder Elbow Surg* 2008;17:e18-20.
20. Craig EV. Subacromial impingement syndrome in hereditary multiple exostoses. *Clin Orthop* 1986;182-4.
21. Fallon PJ, Hollinshead RM. Solitary osteochondroma of the distal clavicle causing a full-thickness rotator cuff tear. *J Shoulder Elbow Surg* 1994;3:266-9.
22. Galanopoulos I, Stavlas P, Beltsios M. Distal Clavicle Osteochondroma Causing Supraspinatus Tendinopathy. *Cureus [Internet]*. [cited 2020 Mar 25];11.
23. Vander Maren C, Guillaumie B, Hüge J et al. [Osteochondroma of the clavicle and pain syndrome of the shoulder. Apropos of a case. Review of the literature]. *Rev Chir Orthop Reparatrice Appar Mot* 1994;80:334-7.
24. Reichmister J, Reeder JD, Gold DL. Osteochondroma of the distal clavicle: an unusual cause of rotator cuff impingement. *Am J Orthop Belle Mead NJ* 2000;29:807-9.
25. Ogawa K, Yoshida A, Ui M. Symptomatic osteochondroma of the clavicle. A report of two cases. *J Bone Joint Surg Am* 1999;81:404-8.
26. Alman BA, Goldberg MJ. Solitary osteochondroma of the clavicle. *J Pediatr Orthop* 1991;11:181-3.
27. Simonetti I, Chianca V, Ascione F et al. Clavicular Osteochondroma: Extremely Rare Cause of Impingement Syndrome. *J Orthop Case Rep* 2018;8:50-3.
28. Al-Qudah AS, Abu-Ali HM, Al-Hussaini MA, Massad IM. Periosteal chondroma of the clavicle: case report and review of the literature. *Int J Surg Lond Engl* 2009;7:140-1.
29. Mollano AV, Hagy ML, Jones KB, Buckwalter JA. Unusual osteochondroma of the medial part of the clavicle causing subclavian vein thrombosis and brachial plexopathy. A case report. *J Bone Joint Surg Am* 2004;86:2747-50.
30. López-Martin N, De Miguel I, Calvo E. Rotator cuff impingement due to enchondroma of the acromion. *Acta Orthop Belg* 2005;71:732-5.
31. Hommadi A, Ziadi T, Louaste J et al. [Chondroma of the acromion: an unusual localization]. *Chir Main* 2009;28:120-2.
32. Çıtlak A, Akgün U, Bulut T, et al. Subacromial osteochondroma: A rare cause of impingement syndrome. *Int J Surg Case Rep* 2015;6C:126-8.
33. Lu MT, Abboud JA. Subacromial osteochondroma. *Orthopedics*. 2011;34:e581-583.
34. Thomas S, Srivastav S, Agarwal S, et al. Arthroscopic excision of subacromial osteochondroma causing impingement of shoulder in a patient with diaphyseal achalasia. *J Clin Orthop Trauma* 2013;4:40-2.
35. Pimpalnerkar A, Anand A, Anand A, et al. Double impingement of the rotator cuff— an unusual presentation of dual osteochondromas. *Eur Orthop Traumatol* 2013;4:169-71.
36. Kumar V, Garg S, Elzein I, et al. Modified Weaver-Dunn procedure versus the use of a synthetic ligament for acromioclavicular joint reconstruction. *J Orthop Surg Hong Kong* 2014;22:199-203.
37. Frost NL, Parada SA, Manoso MW, et al. Scapular osteochondromas treated with surgical excision. *Orthopedics* 2010;33:804.
38. Fukunaga S, Futani H, Yoshiya S. Endoscopically assisted resection of a scapular osteochondroma causing snapping scapula syndrome. *World J Surg Oncol* 2007;5:37.
39. Kumar N, Ramakrishnan V, Johnson GV, Southern S. Endoscopically-assisted excision of scapular osteochondroma. *Acta Orthop Scand* 1999;70:394-6.
40. Rossi B, Fabbriani C, Chalidis BE, et al. Primary malignant clavicular tumours: a clinicopathological analysis of six cases and evaluation of surgical management. *Arch Orthop Trauma Surg* 2011;131:935-9.