

Immune checkpoint inhibitor therapy and outcomes from SARS-CoV-2 infection in patients with cancer: a joint analysis of OnCovid and ESMO-CoCARE registries

Alessio Cortellini ^{1,2}, Gino M Dettorre,³ Urania Dafni,⁴ Juan Aguilar-Company,^{5,6} Luis Castelo-Branco,^{7,8} Matteo Lambertini,^{9,10} Spyridon Gennatas,¹¹ Vasileios Angelis,¹¹ Ailsa Sita-Lumsden,¹² Jacobo Rogado,¹³ Paolo Pedrazzoli,^{14,15} David Viñal,¹⁶ Aleix Prat,^{17,18} Maura Rossi,¹⁹ Rossana Berardi,²⁰ Teresa Alonso-Gordoa,²¹ Salvatore Grisanti,²² Georgia Dimopoulou,⁴ Paola Queirolo,²³ Sylvain Pradervand,²⁴ Alexia Bertuzzi,²⁵ Mark Bower,²⁶ Dirk Arnold,²⁷ Ramon Salazar ²⁸, Marco Tucci,^{29,30} Kevin J Harrington ³¹, Francesca Mazzoni,³² Uma Mukherjee,³³ Zoi Tsourti,³⁴ Olivier Michielin,²⁴ Fanny Pommeret,³⁵ Joan Brunet,³⁶ Bruno Vincenzi,² Giuseppe Tonini,² Andrea Patriarca,³⁷ Federica Biello,³⁸ Marco Krenqli,³⁹ Josep Taberner,⁴⁰ George Pentheroudakis,⁷ Alessandra Gennari,³⁸ Solange Peters,^{7,24} Emanuela Romano,⁴¹ David J Pinato ^{1,38}

To cite: Cortellini A, Dettorre GM, Dafni U, *et al*. Immune checkpoint inhibitor therapy and outcomes from SARS-CoV-2 infection in patients with cancer: a joint analysis of OnCovid and ESMO-CoCARE registries. *Journal for ImmunoTherapy of Cancer* 2022;**10**:e005732. doi:10.1136/jitc-2022-005732

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/jitc-2022-005732>).

ER and DJP contributed equally.

Accepted 17 October 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Alessio Cortellini;
alessiocortellini@gmail.com

ABSTRACT

Background As management and prevention strategies against COVID-19 evolve, it is still uncertain whether prior exposure to immune checkpoint inhibitors (ICIs) affects COVID-19 severity in patients with cancer.

Methods In a joint analysis of ICI recipients from OnCovid (NCT04393974) and European Society for Medical Oncology (ESMO) CoCARE registries, we assessed severity and mortality from SARS-CoV-2 in vaccinated and unvaccinated patients with cancer and explored whether prior immune-related adverse events (irAEs) influenced outcome from COVID-19.

Findings The study population consisted of 240 patients diagnosed with COVID-19 between January 2020 and February 2022 exposed to ICI within 3 months prior to COVID-19 diagnosis, with a 30-day case fatality rate (CFR₃₀) of 23.6% (95% CI 17.8 to 30.7%). Overall, 42 (17.5%) were fully vaccinated prior to COVID-19 and experienced decreased CFR₃₀ (4.8% vs 28.1%, p=0.0009), hospitalization rate (27.5% vs 63.2%, p<0.0001), requirement of oxygen therapy (15.8% vs 41.5%, p=0.0030), COVID-19 complication rate (11.9% vs 34.6%, p=0.0040), with a reduced need for COVID-19-specific therapy (26.3% vs 57.9%, p=0.0004) compared with unvaccinated patients. Inverse probability of treatment weighting (IPTW)-fitted multivariable analysis, following a clustered-robust correction for the data source (OnCovid vs ESMO CoCARE), confirmed that vaccinated patients experienced a decreased risk of death at 30 days (adjusted OR, aOR 0.08, 95% CI 0.01 to 0.69). Overall, 38 patients (15.8%) experienced at least one irAE of any grade at any time prior to COVID-19, at a median

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ SARS-CoV-2 vaccines significantly improve COVID-19 morbidity and mortality in patients with cancer. Efficacy data from large registry studies in patients receiving immune checkpoint inhibitors (ICIs) are still lacking.

WHAT THIS STUDY ADDS

⇒ This joint analysis of patients recently exposed to ICI from OnCovid and European Society for Medical Oncology-CoCARE registries confirms clinical efficacy of SARS-CoV-2 vaccination in reducing COVID-19 morbidity and mortality.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Considering the continuously expanding indication for ICI therapy, these findings are of the utmost importance to ensure effective utilization of this therapy during and beyond the SARS-CoV-2 global pandemic.

time of 3.2 months (range 0.13–48.7) from COVID-19 diagnosis. IrAEs occurred independently of baseline characteristics except for primary tumor (p=0.0373) and were associated with a significantly decreased CFR₃₀ (10.8% vs 26.0%, p=0.0462) additionally confirmed by the IPTW-fitted multivariable analysis (aOR 0.47, 95% CI 0.33 to 0.67). Patients who experienced irAEs also presented

a higher median absolute lymphocyte count at COVID-19 (1.4 vs 0.8 10^9 cells/L, $p=0.0098$).

Conclusion Anti-SARS-CoV-2 vaccination reduces morbidity and mortality from COVID-19 in ICI recipients. History of irAEs might identify patients with pre-existing protection from COVID-19, warranting further investigation of adaptive immune determinants of protection from SARS-CoV-2.

INTRODUCTION

The efficacy of immune checkpoint inhibitors (ICIs) strongly relies on their capacity of inducing T-cell immune reconstitution.¹ T-cell exhaustion is a contributory mechanism underlying the severity of SARS-CoV-2 infection,² leading on one hand to the investigation of programmed-cell death-1 inhibitors as a therapeutic strategy in severe COVID-19.³ On the other hand, given the pathological immune-mediated mechanisms underlying COVID-19 and the risk of immune-pathology stemming from ICI use, there has been growing concern around the use of ICI in patients with COVID-19 and cancer.^{4,5}

Clinical data in support of a protective, as opposed to detrimental, effect of ICI in the prognosis of COVID-19 in patients with cancer have been inevitably biased by patient selection and underlying clinical characteristics. Initial reports revealed inconsistent results ranging from worse outcomes,^{6,7} to no difference in COVID-19 severity^{8,9} in ICI-exposed patients compared with ICI-unexposed patients.

Large meta-analyses have suggested no differential impact of ICIs on COVID-19 morbidity and mortality in comparison to other systemic anticancer therapies.^{10,11}

However, COVID-19 outcomes in patients with cancer have substantially evolved over time. Improved management of COVID-19,¹² immunization campaigns,^{13,14} changes in community transmission and the emergence of new SARS-CoV-2 variants¹⁵ have considerably changed the clinical impact of SARS-CoV-2 infection on patients with cancer since March 2020.

To date, a significant gap in knowledge remains as to whether the positive effect of SARS-CoV-2 vaccination observed in the general population extends to patients with cancer treated with ICI. Recent evidence suggesting that ICI may precipitate subclinical cytokine release following SARS-CoV-2 vaccination¹⁶ strengthens the need to understand the relationship between COVID-19 vaccination and clinical outcomes.

With the aim of providing a contemporary description of COVID-19 morbidity and mortality in patients with cancer who were receiving ICIs at COVID-19 diagnosis and to assess the protective role of SARS-CoV-2 vaccination in this population, we developed this joint analysis of the OnCovid and European Society for Medical Oncology (ESMO) CoCARE registries.

METHODS

Study design

OnCovid (NCT04393974) is a European registry study approved by the UK Health Research Authority (20/HRA/1608) collecting data from consecutive patients fulfilling the following inclusion criteria: (1) age ≥ 18 years; (2) Reverse transcription polymerase chain reaction (RT-PCR) confirmed diagnosis of SARS-CoV-2 infection; (3) history of solid or haematological malignancy either active or in remission at the time of COVID-19 diagnosis.

The ESMO-CoCARE is an observational prospective study, based on a longitudinal multicenter survey of patients with cancer with any solid or hematological malignancy who were diagnosed with COVID-19.

For both registries, data from consecutive, all-comer patients were collected using electronic case report forms designed with the Research Electronic Data Capture software (Vanderbilt University, Nashville, Tennessee, USA). Study details and procedures, patients' eligibility, and clinical endpoints for both studies have already been extensively presented.^{12-14,17-24} A list of participating centers with eligible patients for the present analysis is provided as online supplemental table 1.

Objectives and endpoints

The main objective of this analysis was to assess the protective role of SARS-CoV-2 vaccination in patients with cancer treated with a unique immunotherapy strategy, by comparing COVID-19 morbidity and mortality between unvaccinated and vaccinated patients.

In addition, we aimed to describe differences in COVID-19 severity and mortality depending on prior history of immune-related adverse events (irAEs) captured from ICI initiation until COVID-19 diagnosis.

Data of patients who received ICI within 3 months prior to COVID-19 diagnosis were merged from the OnCovid and ESMO CoCARE registries. Patients on chemotherapy-ICI and targeted therapy-ICI combinations were excluded from the analysis.

To reflect the temporal evolution of the pandemic, we first categorized patients according to date of COVID-19 diagnosis into prevaccination (from February 2020 to November 2020), alpha-delta (B.1.1.7-B.1.617.2) variants (from December 2020 to December 14, 2021), and omicron (B.1.1.529) variant (from December 15, 2021 to February 2022) pandemic phases as previously reported,¹³ and described COVID-19 mortality over time.

All-cause case fatality rate at 30 days (CFR₃₀) was chosen as the main clinical endpoint, to differentiate early COVID-19-related mortality, from late, likely cancer-related deaths. As measures of COVID-19 morbidity, we evaluated the all-cause hospitalization and intensive care unit (ICU) admission rates, the rate of COVID-19 complications (at least one among acute respiratory failure, ARDS, kidney injury, secondary infections, sepsis, septic shock, acute cardiac injury, acute liver injury and others including thrombo-embolic events and other

coagulopathies, autoimmune diseases, gastrointestinal reactions), the receipt of at least one COVID-19-oriented therapy (including antivirals, chloroquine-based treatment, antibiotics, corticosteroids, interleukin-6 inhibitors and others) (yes vs no), and supplemental oxygen therapy requirement (yes vs no).

Patients who received two doses of the BNT162b2, mRNA-1273, ChAdOx1-S, and CoronaVac vaccines prior to COVID-19, or in case of infection diagnosed at least 28 days after a single dose of the Ad.26.COV2.S vaccine, were defined as fully vaccinated. Patients who received one vaccination, without meeting the above-mentioned time criteria, were considered partially vaccinated, while patients who received a third dose of either the BNT162b2 or mRNA-1273 vaccine (or a second dose after the Ad.26.COV2.S vaccine) were considered boosted. Considering the limited sample size of vaccinated patients with breakthrough infections in the study population, and that the electronic case report form of the ESMO-CoCARE registry was not designed to collect information on booster doses, patients were grouped as unvaccinated (including partially vaccinated) and fully vaccinated (either double-dosed or boosted patients) for all the comparative analyses, while patients with unknown vaccination status were excluded.

For the irAEs analysis, we evaluated COVID-19 outcomes according to the experience of any grade (National Cancer Institute Common Toxicity Criteria for Adverse Events, V.5.0) treatment-related side effects with a putative immune-mediated mechanisms at any time prior to COVID-19. These were previously evaluated by clinicians at participating sites during routine consultations as clinically indicated, without predefined time points, and collected retrospectively by investigators.

Considering the recognized role of lymphopenia as prognostic biomarker in patients with COVID-19,²⁵ we explored the association between the absolute lymphocyte count at COVID-19 (within 1 week of diagnosis) and the experience of prior irAEs in the subset of patients from the OnCovid registry. A detailed description of statistical analysis is provided as online supplemental methods.

RESULTS

Study population

By the respective data lock dates of February 4, 2022 and May 17, 2022, the OnCovid and ESMO CoCARE included 3820 and 2310 patients. After the exclusion of ineligible patients, data from 178 (74.2%—OnCovid) and 62 (25.8%—ESMO CoCARE) patients diagnosed with COVID-19 between January 2020 and February 2022, who were receiving ICIs within 3 months prior to SARS-CoV-2 infection diagnosis, were merged.

Figure 1 reports a detailed study flow diagram. The final study population consisted of 240 patients, of whom 130 (54.2%) were diagnosed with COVID-19 during the prevaccination phase, 79 (32.9%) during the alpha–delta phase, and 31 (12.9%) during the omicron phase, with reducing CFR₃₀ over time: 25.8% (24/93 patients, 95% CI 16.5 to 38.4), 31.5% (17/54 patients, 95% CI 18.3 to 50.4), 3.6% (1/28 patients, 95% CI 0.09 to 19.8).

The most frequent primary tumor was lung cancer (47.1%), the majority of patients were male (67.5%), aged ≥65 years (62.1%), with at least one comorbidity (77.1%) and presented an active (76.7%), and advanced-stage (80.2%) tumor (table 1).

The received ICI-based regimens were: 136 (56.7%) PD-1 inhibitors monotherapy, 54 (22.5%) PD-L1 inhibitors monotherapy, 20 (8.3%) others/experimental




Figure 1 Study flow diagram. ESMO, European Society for Medical Oncology; ICI, immune checkpoint inhibitor; SACT, systemic anti-cancer therapies.

Table 1 Baseline patient characteristic and COVID-19 outcomes of the study population

	ICI population N=240 (%)
Country	
UK	47 (19.6)
Spain	77 (32.1)
Italy	90 (37.5)
Others	26 (10.8)
Sex	
Female	78 (32.5)
Male	162 (67.5)
Age	
<65 years	90 (37.5)
≥65 years	149 (62.1)
Missing	1 (0.4)
Comorbidities	
No	55 (22.9)
Yes	185 (77.1)
Primary tumor	
Lung	113 (47.1)
Melanoma	51 (21.2)
Others	76 (31.7)
Tumor stage	
Non-advanced	37 (15.6)
Advanced	190 (80.2)
Missing	10 (4.2)
Tumor status at COVID-19 diagnosis	
Remission/in-response	52 (21.7)
Active malignancy	184 (76.7)
Missing	4 (1.7)
SARS-CoV-2 vaccination status	
Unvaccinated	182 (75.8)
Fully vaccinated	42 (17.5)
Partially vaccinated	3 (1.3)
Unkown	13 (5.4)
COVID-19 outcomes	
	N (rate, 95% CI)
Oxygen therapy	81 (37.6 , 29.9 to 46.8)
Missing	25
COVID-19-specific therapy	114 (51.6 , 42.5 to 61.9)
Missing	19
Complications from COVID-19	73 (30.4 , 23.8 to 38.2)
Hospitalization	131 (56.2 , 47.0 to 66.7)
Missing	7
ICU admission	22 (9.4 , 5.9 to 14.3)
Missing	7
30-days case fatality rate	55 (23.6 , 17.8 to 30.7)
Missing	7

COVID-19 outcomes' rates are provided in bold.
ICI, immune checkpoint inhibitor; ICU, intensive care unit.

ICIs, 19 (7.9%) CTLA-4/PD-1 inhibitors combinations and 11 (4.6%) not specified chemotherapy-free ICI regimens.

Most patients were unvaccinated prior to COVID-19 (75.8%), 17.5% were fully vaccinated, 1.3% partially vaccinated, while vaccination status was unknown for 13 patients (5.4%). Among fully vaccinated patients, 17 from the OnCovid registry received a booster dose. Vaccination details for both the registries are summarized in online supplemental table 2.

The median observation period for the whole cohort was 91 days (IQR: 15.8–319.0) and the CFR₃₀ was 23.6% (95% CI 17.8% to 30.7%). All COVID-19 outcomes for the whole cohort are summarized in [table 1](#)

SARS-CoV-2 vaccination is associated with improvement in COVID-19 outcomes in ICI recipients

After the exclusion of 13 patients with unknown vaccination status, 227 patients were included in the SARS-CoV-2 vaccine analysis.

None of the baseline demographics and oncological characteristics were associated with SARS-CoV-2 vaccination status, with the exception of a higher proportion of patients with at least one comorbidity among unvaccinated patients (80.5% vs 64.3%, $p=0.0230$) (online supplemental table 3).

Univariable analysis revealed that fully vaccinated patients experienced decreased rates of death at 30 days (4.8% vs 28.1%, $p=0.0009$), hospitalization (27.5% vs 63.2%, $p<0.0001$), COVID-19 complications (11.9% vs 34.6%, $p=0.0040$), reduced need for COVID-19-specific therapy (26.3% vs 57.9%, $p=0.0004$) and oxygen therapy (15.8% vs 41.5%, $p=0.0030$) in comparison to unvaccinated/partially vaccinated patients. We found no significant difference in terms of ICU admission rates, despite arithmetically fewer vaccinated patients being admitted to ICU (4.8% vs 28.1%, $p=0.14$) ([figure 2](#), online supplemental table 4).

Distribution of baseline patient characteristics prior to and after inverse probability of treatment weighting (IPTW) is reported in online supplemental table 5. Given the suboptimal balancing ability, country, comorbidities, tumor status and tumor stage were included in all IPTW-fitted multivariable logistic regression models for each COVID-19 outcome, which are reported in full as online supplemental table 6 and are summarized in the forest plot graph provided in [figure 3](#). Compared with unvaccinated patients, full vaccination was associated with a decreased risk of death at 30 days (adjusted OR, aOR 0.08, 95% CI 0.03 to 0.26), of hospitalization (aOR 0.15, 95% CI 0.07 to 0.36), of COVID-19 complications (aOR 0.24, 95% CI 0.12 to 0.49) and of need for COVID-19-specific therapy (aOR 0.25, 95% CI 0.13 to 0.46). However, after clustered-robust correction for data source, the upper limit CI crosses one for all COVID-19 outcomes except for CFR₃₀ (aOR 0.08, 95% CI 0.01 to 0.69).

History of irAEs prior to COVID-19 is associated with decreased COVID-19 mortality in patients receiving ICI

Overall, 38 patients (15.8%) experienced any grade irAEs at any time prior to COVID-19, which are summarized in online supplemental table 7. The median time from occurrence of irAEs and COVID-19 diagnosis was 3.2 months (range 0.13–48.7, computed on data of 27 patients from the OnCovid registry).

The occurrence of irAEs was not associated with any of the baseline demographics and oncological characteristics, including the disease status (active vs remissive/in response) at COVID-19 ($p=0.5339$), with the exception of the primary tumor ($p=0.0373$) (online supplemental table 8).

Univariable analysis showed similar rates of hospitalization (51.3% vs 57.1%, $p=0.5158$), ICU admission (16.2% vs 8.1%, $p=0.1252$), COVID-19 complications (23.7% vs 31.7%, $p=0.3265$), COVID-19-specific therapy (45.7% vs 52.6%, $p=0.4498$) and oxygen requirement (39.3% vs 37.4%, $p=0.8251$) between patients who experienced and those who did not experience irAEs prior to COVID-19 (online supplemental table 9). However, the occurrence of irAEs was associated with a significantly decreased CFR₃₀ (10.8% vs 26.0%, $p=0.0462$) ([figure 4A](#)).

Distribution of baseline characteristics distribution prior to and after the IPTW is reported in online supplemental table 10. Given the suboptimal balancing ability, country, tumor stage, primary tumor and vaccination status were included in the IPTW-fitted multivariable logistic regression model for COVID-19 mortality, which confirmed that patients who experienced any grade irAEs prior to COVID-19 had a decreased risk of death at 30 days (aOR 0.47, 95% CI 0.23 to 0.99). Clustered-robust correction for data source further strengthened this finding (aOR 0.47, 95% CI 0.33 to 0.67) (online supplemental table 11).

Lastly, in a subset of patients from the OnCovid cohort, we revealed that the median absolute lymphocyte count within 1 week of COVID-19 diagnosis was significantly higher among patients who experienced any grade irAEs prior to COVID-19 than in those who did not experience irAEs (1.4 vs 0.8 10^9 cells/L, $p=0.0098$) ([figure 4B](#)).

DISCUSSION

Our study is the largest analysis on patients with cancer on ICIs diagnosed with COVID-19 to date. With the inclusion of patients diagnosed up until February 2022, it provides a more contemporary picture of COVID-19 outcomes in this specific population. Although merely descriptive due to the limited sample size of subgroups, the reducing CFR₃₀ across the pandemic phases suggests a time-dependent improvement of COVID-19 mortality, especially during the more recent Omicron outbreak, as already reported for the OnCovid population.¹³

Even considering the time requirements for the delivery of immunization campaigns since the first SARS-CoV-2 vaccine approval,²⁶ and that most of the included

COVID-19 outcomes according to the SARS-CoV-2 vaccination status




Figure 2 Histogram plot summarizing all COVID-19 outcomes according to the vaccination status. All rates with 95% CIs are available in online supplemental table 4. CFR₃₀, 30-day case fatality rate; ICU, intensive care unit.

patients were diagnosed during the prevaccination phase, we consider 17.5% of full vaccination a relatively low rate, and a possible impact of vaccine hesitancy, as initially reported in early 2021,^{27 28} cannot be excluded.

Although preliminary evidence from clinical trials supports the safety and immunogenicity of SARS-CoV-2 vaccines in patients with cancer on active ICI-based treatments,^{16 29 30} this study demonstrates the efficacy of anti-COVID-19 vaccination in patients receiving ICI in routine clinical practice. The ~83% reduction in the CFR₃₀ in fully vaccinated patients along with COVID-19-related morbidity is confirmed after adjustment for major prognostic confounders in IPTW-fitted models, a process made necessary by the inherent differences existing in study procedures and data collection modalities between the two registries.

The convergence of COVID-19 and ICI-toxicity in eliciting unopposed T-cell activation and downstream cytokine excess has been highlighted suggested as a hypothetical source of clinical risk to patients with cancer ever since the beginning of the pandemic.^{5 31} Contrary to initial concerns, we document an association between the occurrence of irAEs and reduced CFR₃₀; a novel finding of potential interest in the development of COVID-19-specific therapeutics.

In our study, the protective role of irAEs of all grades on COVID-19-related mortality was independent of common clinicopathological features relating to cancer and COVID-19 prognosis, including SARS-CoV-2 vaccination status. It has been established that patients experiencing irAEs are those capable of mounting a more vigorous anticancer immune reconstitution, resulting in

IPTW fitted multivariable analysis of COVID-19 outcomes




Figure 3 Summary of the inverse probability of treatment weighing (IPTW) fitted multivariable analyses for each COVID-19 outcomes according to the vaccination status prior to (blue) and after (red) the clustered-robust SE and 95% CI adjustments for the data source. Adjusting covariates for each COVID-19 outcome were country of origin, comorbidities, tumor status, and tumor stage at COVID-19. Full multivariable models are available in online supplemental table 6. aOR, adjusted OR; CFR30, 30-day case fatality rate.



A All-cause CFR₃₀ according to irAEs experience

B


Figure 4 (A) Histogram plot summarizing the all-cause 30-day case fatality rate (CFR30) analysis according to the occurrence of any grade immune-related adverse events prior to COVID-19. Inverse probability of treatment weighing (IPTW) fitted adjusted OR for the risk of death at 30 days with clustered robust 95% CI correction for the data source is presented. All rates with 95% CI are available in online supplemental table 9. Adjusting covariates were country of origin, primary tumor, tumor stage at COVID-19 and vaccination status. Full multivariable model is available in online supplemental table 11. (B) Violin plot reporting the median absolute lymphocyte count at COVID-19 (within 1 week of diagnosis) according to the prior occurrence or any grade irAEs. irAEs, immune-related adverse events.

longer survival.³² Because T-cell exhaustion is not solely a hallmark of cancer progression but a mechanism of COVID-19 severity,^{25 31} we speculate whether history of prior irAE might be a surrogate of more functional T-cell immunity, leading to improved mortality from COVID-19 irrespective of vaccine status.

In keeping with this view, we found that the absolute lymphocyte count at COVID-19, was significantly higher among patients who experienced prior irAEs. It is well known that patients with severe COVID-19 show reduced counts of peripheral CD4+andCD8+ T cells³¹, and that reduced CD4+/CD8+T cells, B cells, NK cells, and absolute lymphocyte cell count levels are significantly associated with COVID-19 mortality in the general population.²⁵ At the same time, the known mechanisms leading to irAEs involve expansion of intratumoral and peripheral T-cell receptor repertoires along with a mobilization of large numbers of T cells^{33 34} and, to a lesser extent, activation and exhaustion of CD21^{low} B cells.³⁵ On the other hand, a decrease in the absolute lymphocyte count has been reported with severe ICI-associated myocarditis.³⁶

While OnCovid and ESMO CoCARE registries lack information on T-cell phenotype at COVID-19 diagnosis, our findings are provocative in suggesting that prior irAE might represent a hallmark of protection from COVID-19 mortality through invigorated T-cell immunity. These findings deserve further mechanistic studies to fully elucidate the immunological links between irAEs and COVID-19 outcomes in patients with cancer.

Our study acknowledges several limitations, including lack of data regarding the smoking status and more detailed information regarding irAEs duration and management. Of note, previous irAEs and their putative immune-mediated mechanism were assessed at participating site in routine practice, without predefined time points. This might have impacted the quality of data with risks of underreporting, as the 16.7% and 3.1% rates of all grade and \geq G3 irAEs, respectively, are lower than those reported in interventional clinical trials with ICI-based regimens,³⁷ but comparable to reports from clinical practice.³⁸

In addition, inherent differences between the two registries significantly impacted the accuracy of the estimates from the vaccination analysis: information about booster doses only recently started to be collected for patients entered in the ESMO CoCARE registry and was not available for our analysis. Furthermore, for ~24% of vaccinated patients, the specific type of vaccine could not be reconstructed. While constituting an important limitation, this is unlikely to have affected our results, given recent evidence suggesting largely comparable efficacy of commonly available SARS-CoV-2 vaccines.³⁹

Lastly, despite the inclusion of a significant proportion of more recently diagnosed patients, the lack of availability of viral genomic sequences across the pandemic phases did not allow us to make conclusive considerations about new SARS-CoV-2 variants, while the limited sample size of the ‘alpha–delta’ and ‘omicron’ phases subgroups

prevented us from running adequately powered time-adjusted analyses.

Despite the mentioned limitations, our results collectively support the notion that ICI recipients are not especially vulnerable to COVID-19, with mortality rates that are in keeping with the general population with COVID-19 and cancer. In these patients, SARS-CoV-2 vaccination leads to significantly improved outcome from COVID-19, comparably with other oncological patient populations.^{13 14 40} Considering the continuously expanding indication for ICI therapy,⁴¹ our findings are of utmost importance to ensure effective utilization of this therapy during and beyond the SARS-CoV-2 global epidemic.

Author affiliations

¹Department of Surgery & Cancer, Hammersmith Hospital Campus, Imperial College London, London, UK

²Medical Oncology, Fondazione Policlinico Universitario Campus Bio-Medico, Via Alvaro del Portillo, 200 - 00128, Roma, Italy

³Department of Internal Medicine, Washington University School of Medicine, St. Louis, Missouri, USA

⁴Laboratory of Biostatistics, School of Health Sciences, National and Kapodistrian University of Athens, Athens, Greece

⁵Medical Oncology, Vall d'Hebron University Hospital and Institute of Oncology (VHIO), Barcelona, Spain

⁶Infectious Disease, Vall d'Hebron University Hospital, Barcelona, Spain

⁷Scientific and Medical Division, ESMO (European Society for Medical Oncology), Lugano, Switzerland

⁸NOVA National School of Public Health, NOVA University, Lisbon, Portugal

⁹Department of Internal Medicine and Medical Specialties (DiMI), School of Medicine, University of Genoa, Genoa, Italy

¹⁰Medical Oncology Department, U.O. Clinica di Oncologia Medica, IRCCS Ospedale Policlinico San Martino, Genova, Italy

¹¹Medical Oncology Department, The Royal Marsden Hospital and NHS Foundation Trust, London, UK

¹²Medical Oncology, Guy's and St Thomas' NHS Foundation Trust (GSTT), London, UK

¹³Medical Oncology Department, Hospital Universitario Infanta Leonor, Madrid, Spain

¹⁴Medical Oncology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

¹⁵Department of Internal Medicine and Medical Therapy, University of Pavia, Pavia, Italy

¹⁶Medical Oncology Department, Hospital Universitario La Paz, Madrid, Spain

¹⁷Department of Medical Oncology, Hospital Clinic de Barcelona, Barcelona, Spain

¹⁸Translational Genomics and Targeted Therapies in Solid Tumors, Institut d'Investigacions Biomediques August Pi i Sunyer (IDIBAPS), Barcelona, Spain

¹⁹Oncology Department, ASO 'SS Antonio Biagio e Cesare Arrigo', Alessandria, Italy

²⁰Medical Oncology, AOU Ospedali Riuniti, Polytechnic University of the Marche Region, Ancona, Italy

²¹Medical Oncology Department, Hospital Universitario Ramón y Cajal, Madrid, Spain

²²Medical Oncology Unit, Spedali Civili, Brescia, Italy

²³Melanoma Sarcoma and Rare Tumors, IEO, European Institute of Oncology IRCCS, Milan, Italy

²⁴Oncology Department, Centre Hospitalier Universitaire Vaudois (CHUV), Lausanne, Switzerland

²⁵Medical Oncology and Hematology Unit, Humanitas Cancer Center, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy

²⁶Department of Oncology and National Centre for HIV Malignancy, Chelsea and Westminster Hospital, London, UK

²⁷Oncology, Haematology, Palliative Care Department, Asklepios Klinik Altona e Asklepios Kliniken, Hamburg, Germany

²⁸Department of Medical Oncology, ICO L'Hospitalet, Oncobell Program (IDIBELL), CIBERONC, Hospitalet de Llobregat, Barcelona, Spain

²⁹Section of Medical Oncology, Department of Interdisciplinary Medicine (DIM), University of Bari 'Aldo Moro', Bari, Italy

³⁰IRCCS Istituto Tumori Giovanni Paolo II, Bari, Italy

- ³¹Division of Radiotherapy and Imaging, The Royal Marsden Hospital and The Institute of Cancer Research NIHR Biomedical Research Centre, London, UK
- ³²Medical Oncology, Careggi University Hospital, Florence, Italy
- ³³Medical Oncology, Barts Health NHS Trust, London, UK
- ³⁴Frontier Science Foundation-Hellas, Athens, Greece
- ³⁵Department of Cancer Medicine, Institut Gustave Roussy, University of Paris Saclay, 114 rue Edouard Vaillant, Villejuif, France
- ³⁶Department of Medical Oncology, Catalan Institute of Oncology, University Hospital Josep Trueta, Girona, Spain
- ³⁷Division of Haematology, Department of Translational Medicine, University of Piemonte Orientale and Azienda Ospedaliera Maggiore della Carità, Novara, Italy
- ³⁸Division of Oncology, Department of Translational Medicine, University of Piemonte Orientale and Azienda Ospedaliera Maggiore della Carità, Novara, Italy
- ³⁹Division of Radiotherapy, Department of Translational Medicine, University of Piemonte Orientale and Azienda Ospedaliera Maggiore Della Carita, Novara, Italy
- ⁴⁰Medical Oncology, Vall d'Hebron University Hospital and Institute of Oncology (VHIO), IOB-Quiron, UVic-UCC, Barcelona, Spain
- ⁴¹Center for Cancer Immunotherapy, Department of Oncology, PSL Research University, Institut Curie, Paris, France

Twitter Gino M Dettorre @DettorreGino, Matteo Lambertini @matteolambe, Aleix Prat @prat_aleix and Ramon Salazar @RamonSalazarS

Contributors All authors contributed to the publication according to the ICMJE guidelines for the authorship. All authors had access to all the data reported in the study. All authors read and approved the submitted version of the manuscript (and any substantially modified version that involves the author's contribution to the study). Each author has agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work, even ones in which the author was not personally involved, are appropriately investigated, resolved, and the resolution documented in the literature. The corresponding author (AC) had full access to all of the data and acts as guarantor of the overall content of the study. AC accepts full responsibility for the work and/or the conduct of the study, had access to the data, and controlled the decision to publish. AC, DJP, SP, ZT, and UD have accessed and verified the study data. Study concept and design: AC, ER, and DJP. Acquisition of data: AC, GMD, UD, JA-C, LC-B, ML, SG, VA, AS-L, JR, PP, DV, AP, MR, RB, TA-G, SG, GD, PQ, SP, AB, MB, DA, RS, MT, KJH, FM, UM, ZT, OM, FP, JB, BV, AP, FB, MK, JT, GP, AG, SP, ER, and DJP. Analysis and interpretation of data: AC, ER, DJP. Drafting of the manuscript: AC, ER, DJP. Statistical analysis: AC, ZT, and UD. Manuscript review and approval: AC, GMD, UD, JA-C, LC-B, ML, SG, VA, AS-L, JR, PP, DV, AP, MR, RB, TA-G, SG, GD, PQ, SP, AB, MB, DA, RS, MT, KJH, FM, UM, ZT, OM, FP, JB, BV, AP, FB, MK, JT, GP, AG, SP, ER, and DJP. Obtained funding: DJP and ER. Study supervision: ER and DJP.

Funding OnCovid is sponsored by Imperial College London and received direct project funding and infrastructural support by the NIHR Imperial Biomedical Research Centre (BRC).

Disclaimer The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. Neither sponsor nor the funders of the study had any role in study design, data collection, data analysis, data interpretation, or writing of the report. ESMO CoCARE is supported by a grant from the European Society of Medical Oncology.

Competing interests KJH declares research funding from AstraZeneca, Boehringer-Ingelheim, MSD, Replimune and advisory board fees/honoraria from Arch Oncology, AstraZeneca, BMS, Boehringer-Ingelheim, Codiak Biosciences, Inzen Therapeutics, Merck-Serono, MSD, Pfizer, Replimune. DA reports consultation/advisory role for AstraZeneca, Bristol Myers Squibb, Merck Sharp reports speaker's engagement from AstraZeneca, Bristol Myers Squibb, Merck Sharp reports serving as local PI for Bristol Myers Squibb, Pierre Fabre Pharma and coordinating PI for OncoLytics; reports grant funding from AbbVie; reports being/been DSMB chair of Sanofi (Genzyme); reports being/been a steering committee member of Roche. Olivier Michielin reports personal fees from Bristol-Myers Squibb, MSD, Novartis, Roche, Amgen, NeraCare, outside the submitted work. JR received speaker or advisory fees from Roche, Astra Zeneca, Merck, Ferrer, Persan Farma, Teva Pharma, Leo Pharma, Fresenius kabi, MSD, BMS. Travel expenses support from BMS, MSD, RocheUrania Dafni reports honorarium as Member of the Tumor Agnostic Evidence Generation Working Group of Roche, outside the submitted work. GP reports grants from Amgen, Lilly; grants, personal fees and nonfinancial support from Merck; grants and non-financial support from AstraZeneca; grants and personal fees from Roche, Bristol Myers Squibb, MSD, Novartis, outside the submitted work. Solange

Peters reports consultation/advisory role for AbbVie, Amgen, AstraZeneca, Bayer, BeiGene, Biocartis, Bio Invent, Blueprint Medicines, Boehringer Ingelheim, Bristol-Myers Squibb, Clovis, Daiichi Sankyo, Debiopharm, Eli Lilly, Elsevier, F Hoffmann-La Roche/Genentech, Foundation Medicine, Illumina, Incyte, IQVIA, Janssen, Medscape, Merck Sharp and Dohme, Merck Serono, Merrimack, Mirati, Novartis, Pharma Mar, Phosphatin Therapeutics, Pfizer, Regeneron, Sanofi, Seattle Genetics, Takeda, Vaccibody; talk in a company's organized public event for AstraZeneca, Boehringer Ingelheim, Bristol-Myers Squibb, e-cancer, Eli Lilly, F. Hoffmann-La Roche/Genentech, Illumina, Medscape, Merck Sharp and Dohme, Novartis, PER, Pfizer, Prime, RTP, Sanofi, Takeda; receipt of grants/research supports from being a (sub) investigator in trials (institutional financial support for clinical trials) sponsored by Amgen, AstraZeneca, Biodesix, Boehringer Ingelheim, Bristol-Myers Squibb, Clovis, F Hoffmann-La Roche/Genentech, GSK, Illumina, Lilly, Merck Sharp and Dohme, Merck Serono, Mirati, Novartis, and Pfizer, Phosphatin Therapeutics. Emanuela Romano reports investigator-initiated trial (funds paid to the institution) supported by Astra-Zeneca, BMS; serves on the consultancy/advisory board for Astra-Zeneca, Merck, Roche, Pierre Fabre. ML acted as consultant for Roche, Novartis, Lilly, AstraZeneca, Exact Sciences, MSD, Pfizer, Seagen and received speaker honoraria from Roche, Novartis, Lilly, Pfizer, Takeda, Ipsen and Sandoz outside the submitted work. Alessandra Gennari has declared consulting/advisory role for Roche, MSD, Eli Lilly, Pierre Fabre, Eisai, and Daiichi Sankyo; speakers bureau for Eisai, Novartis, Eli Lilly, Roche, Teva, Gentili, Pfizer, Astra Zeneca, Celgene, and Daiichi Sankyo; research funds: Eisai, Eli Lilly, and Roche. CMV has received travel grants and other honoraria from BMS, MSD, Novartis and Roche. Joan Brunet has declared consulting/advisory role for MSD and Astra Zeneca. Aleix Prat has declared personal honoraria from Pfizer, Roche, MSD Oncology, Eli Lilly, and Daiichi Sankyo; travel, accommodations, and expenses paid by Daiichi Sankyo; research funding from Roche and Novartis; and consulting/advisory role for NanoString Technologies, Amgen, Roche, Novartis, Pfizer and Bristol-Myers Squibb. Mark Bower received speakers' fee from Eisai pharma, Gilead Sciences, Merck and Viiv. JT reports consulting fees from Array Biopharma, AstraZeneca, Avvinity, Bayer, Boehringer Ingelheim, Chugai, Daiichi Sankyo, F Hoffmann-La Roche, Genentech, HalioDX SAS, Hutchison MediPharma International, Ikena Oncology, Inspira, IQVIA, Lilly, Menarini, Merck Serono, Merus, MSD, Mirati, Neophore, Novartis, Ona Therapeutics, Orion Biotechnology, Peptomyc, Pfizer, Pierre Fabre, Samsung Bioepis, Sanofi, Seattle Genetics, Scandion Oncology, Servier, Sotio Biotech, Taiho, Tessa Therapeutics, and TheraMyc; speaker's fees from Imedex, Medscape Education, MJH Life Sciences, PeerView Institute for Medical Education, and Physicians Education Resource; and institutional research support from Amgen, Array Biopharma, AstraZeneca Pharmaceuticals, BeiGene, Boehringer Ingelheim, Bristol Myers Squibb, Celgene, Debiopharm International, F Hoffmann-La Roche, Genentech, HalioDX, Hutchison MediPharma International, Janssen-Cilag, MedImmune, Menarini, Merck Health, Merck Sharp and has had leadership roles in the European AIDS Clinical Society, UNAIDS, WHO, and The European Hematology Association/ European Society of Medical Oncology. DJP received lecture fees from Viiv Healthcare, Bayer Healthcare, BMS, Roche, Eisai, Falk Foundation, travel expenses from BMS and Bayer Healthcare; consulting fees from Mina Therapeutics, Eisai, Roche, DaVolterra and Astra Zeneca; research funding (to institution) from MSD and BMS. AC received consulting fees from MSD, BMS, AstraZeneca, Roche; speakers' fee from AstraZeneca, MSD, Novartis and Eisai. All remaining authors have declared no conflicts of interest.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Individual, deidentified participant data and data dictionary may be made available at the request of investigators whose proposed use of the data has been approved by the OnCovid consortium and ESMO CoCARE steering committees following review of a methodologically sound research proposal.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any

purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Alessio Cortellini <http://orcid.org/0000-0002-1209-5735>

Ramon Salazar <http://orcid.org/0000-0001-9419-6232>

Kevin J Harrington <http://orcid.org/0000-0002-6014-348X>

David J Pinato <http://orcid.org/0000-0002-3529-0103>

REFERENCES

- 1 Bagchi S, Yuan R, Engleman EG. Immune checkpoint inhibitors for the treatment of cancer: clinical impact and mechanisms of response and resistance. *Annu Rev Pathol* 2021;16:223–49.
- 2 Kusnadi A, Ramirez-Suástegui C, Fajardo V, et al. Severely ill COVID-19 patients display impaired exhaustion features in SARS-CoV-2-reactive CD8⁺ T cells. *Sci Immunol* 2021;6:eabe4782.
- 3 Awadasseid A, Yin Q, Wu Y, et al. Potential protective role of the anti-PD-1 blockade against SARS-CoV-2 infection. *Biomed Pharmacother* 2021;142:111957.
- 4 Garassino MC, Ribas A. At the crossroads: COVID-19 and Immune-Checkpoint blockade for cancer. *Cancer Immunol Res* 2021;9:261–4.
- 5 Bersanelli M. Controversies about COVID-19 and anticancer treatment with immune checkpoint inhibitors. *Immunotherapy* 2020;12:269–73.
- 6 Bersanelli M, Giannarelli D, De Giorgi U, et al. Symptomatic COVID-19 in advanced-cancer patients treated with immune-checkpoint inhibitors: prospective analysis from a multicentre observational trial by FICOG. *Ther Adv Med Oncol* 2020;12:1758835920968463.
- 7 Robilotti EV, Babady NE, Mead PA, et al. Determinants of COVID-19 disease severity in patients with cancer. *Nat Med* 2020;26:1218–23.
- 8 Luo J, Rizvi H, Egger JV, et al. Impact of PD-1 blockade on severity of COVID-19 in patients with lung cancers. *Cancer Discov* 2020;10:1121–8.
- 9 Rogiers A, Pires da Silva I, Tentori C, et al. Clinical impact of COVID-19 on patients with cancer treated with immune checkpoint inhibition. *J Immunother Cancer* 2021;9:e001931.
- 10 Liu Y, Liu S, Qin Y, et al. Does prior exposure to immune checkpoint inhibitors treatment affect incidence and mortality of COVID-19 among the cancer patients: the systematic review and meta-analysis. *Int Immunopharmacol* 2021;101:108242.
- 11 Lazarus G, Budiman RA, Rinaldi I. Does immune checkpoint inhibitor increase the risks of poor outcomes in COVID-19-infected cancer patients? A systematic review and meta-analysis. *Cancer Immunol Immunother* 2022;71:373–86.
- 12 Pinato DJ, Patel M, et al, OnCovid Study Group. Time-Dependent COVID-19 mortality in patients with cancer: an updated analysis of the OnCovid registry. *JAMA Oncol* 2022;8:114–22.
- 13 Pinato DJ, Aguilar-Company J, Ferrante D. Outcomes of the SARS-CoV-2 omicron (B.1.1.529) variant outbreak among vaccinated and unvaccinated patients with cancer in Europe: results from the retrospective, multicentre, OnCovid registry study. *Lancet Oncol* 2022.
- 14 Pinato DJ, Ferrante D, Aguilar-Company J, et al. Vaccination against SARS-CoV-2 protects from morbidity, mortality and sequelae from COVID-19 in patients with cancer. *Eur J Cancer* 2022;171:64–74.
- 15 Callaway E. Beyond omicron: what's next for COVID's viral evolution. *Nature* 2021;600:204–7.
- 16 Walle T, Bajaj S, Kraske JA, et al. Cytokine release syndrome-like serum responses after COVID-19 vaccination are frequent and clinically inapparent under cancer immunotherapy. *Nat Cancer* 2022;3:1039–51.
- 17 Pinato DJ, Lee AJX, Biello F, et al. Presenting features and early mortality from SARS-CoV-2 infection in cancer patients during the initial stage of the COVID-19 pandemic in Europe. *Cancers* 2020;12. doi:10.3390/cancers12071841. [Epub ahead of print: 08 07 2020].
- 18 Pinato DJ, Scotti L, Gennari A, et al. Determinants of enhanced vulnerability to coronavirus disease 2019 in UK patients with cancer: a European study. *Eur J Cancer* 2021;150:190–202.
- 19 Pinato DJ, Tabernero J, Bower M, et al. Prevalence and impact of COVID-19 sequelae on treatment and survival of patients with cancer who recovered from SARS-CoV-2 infection: evidence from the OnCovid retrospective, multicentre registry study. *Lancet Oncol* 2021;22:1669–80.
- 20 Pinato DJ, Zambelli A, Aguilar-Company J, et al. Clinical portrait of the SARS-CoV-2 epidemic in European cancer patients. *Cancer Discov* 2020. doi:10.1158/2159-8290.CD-20-0773. [Epub ahead of print: 31 Jul 2020].
- 21 Cortellini A, Salazar R, Gennari A, et al. Persistence of long-term COVID-19 sequelae in patients with cancer: an analysis from the OnCovid registry. *Eur J Cancer* 2022;170:10–16.
- 22 Cortellini A, Gennari A, Pommeret F, et al. COVID-19 sequelae and the host proinflammatory response: an analysis from the OnCovid registry. *J Natl Cancer Inst* 2022;114:979–87.
- 23 ESMO-COCARE registry. Available: <https://www.esmo.org/covid-19-and-cancer/registries-studies-and-surveys/esmo-cocare-registry> [Accessed 26 Jun 2022].
- 24 Castelo-Branco L, Tsourti Z, Gennatas S, et al. COVID-19 in patients with cancer: first report of the ESMO international, registry-based, cohort study (ESMO-CoCARE). *ESMO Open* 2022;7:100499.
- 25 Huang W, Berube J, McNamara M, et al. Lymphocyte subset counts in COVID-19 patients: a meta-analysis. *Cytometry A* 2020;97:772–6.
- 26 Covid-19 vaccine: first person receives pfizer Jab in UK. Available: <https://www.bbc.com/news/uk-55227325> [Accessed 23 Aug 2022].
- 27 Ingram SA, Caston NE, Andrews CJ, et al. Hesitancy and malignancy: vaccine hesitancy among individuals with cancer. *Journal of Clinical Oncology* 2021;39:148.
- 28 Villarreal-Garza C, Vaca-Cartagena BF, Becerril-Gaitan A, et al. Attitudes and factors associated with COVID-19 vaccine Hesitancy among patients with breast cancer. *JAMA Oncol* 2021;7:1242–4.
- 29 Waissengrin B, Agbarya A, Safadi E, et al. Short-term safety of the BNT162b2 mRNA COVID-19 vaccine in patients with cancer treated with immune checkpoint inhibitors. *Lancet Oncol* 2021;22:581–3.
- 30 Hibino M, Uryu K, Takeda T, et al. Safety and immunogenicity of mRNA vaccines against severe acute respiratory syndrome coronavirus 2 in patients with lung cancer receiving immune checkpoint inhibitors: a multicenter observational study in Japan. *J Thorac Oncol* 2022;17:1002–13.
- 31 Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med* 2020;8:420–2.
- 32 Cortellini A, Buti S, Agostinelli V, et al. A systematic review on the emerging association between the occurrence of immune-related adverse events and clinical outcomes with checkpoint inhibitors in advanced cancer patients. *Semin Oncol* 2019;46:362–71.
- 33 Weinmann SC, Pisetsky DS. Mechanisms of immune-related adverse events during the treatment of cancer with immune checkpoint inhibitors. *Rheumatology* 2019;58:vii59–67.
- 34 Lozano AX, Chaudhuri AA, Nene A, et al. T cell characteristics associated with toxicity to immune checkpoint blockade in patients with melanoma. *Nat Med* 2022;28:353–62.
- 35 Das R, Bar N, Ferreira M, et al. Early B cell changes predict autoimmunity following combination immune checkpoint blockade. *J Clin Invest* 2018;128:715–20.
- 36 Drobni ZD, Zafar A, Zubiri L, et al. Decreased absolute lymphocyte count and increased neutrophil/lymphocyte ratio with immune checkpoint inhibitor-associated myocarditis. *J Am Heart Assoc* 2020;9:e018306.
- 37 Arnaud-Coffin P, Maillet D, Gan HK, et al. A systematic review of adverse events in randomized trials assessing immune checkpoint inhibitors. *Int J Cancer* 2019;145:639–48.
- 38 Raschi E, Gatti M, Gelsomino F, et al. Lessons to be learnt from real-world studies on immune-related adverse events with checkpoint inhibitors: a clinical perspective from pharmacovigilance. *Target Oncol* 2020;15:449–66.
- 39 Munro APS, Janani L, Cornelius V, et al. Safety and immunogenicity of seven COVID-19 vaccines as a third dose (booster) following two doses of ChAdOx1 nCov-19 or BNT162b2 in the UK (COV-BOOST): a blinded, multicentre, randomised, controlled, phase 2 trial. *Lancet* 2021;398:2258–76.
- 40 Bestvina CM, Whisenant JG, Torri V, et al. Coronavirus disease 2019 outcomes, patient vaccination status, and cancer-related delays during the omicron wave: a brief report from the TERA-VOLT analysis. *JTO Clin Res Rep* 2022;3:100335.
- 41 Twomey JD, Zhang B. Cancer immunotherapy update: FDA-approved checkpoint inhibitors and companion diagnostics. *Aaps J* 2021;23:39.