

Twist-induced orbital chirality in a photonic laser

Kristina Frizyuk & Costantino De Angelis

Check for updates

The combination of two achiral photonic membranes into a twisted bilayer breaks mirror symmetry and induces non-Hermitian coupling between counter-rotating collective modes, making laser emission chiral.

In light-matter interactions, chirality is traditionally associated with circular polarization or geometrical characteristics of the system. Generating chiral laser emission therefore usually relies on the excitation of circularly polarized modes, using chiral cavities or introducing asymmetric excitation^{1–3}. Now writing in *Nature Communications*, Peng and co-workers demonstrate a fundamentally different mechanism to achieve chiral lasing, where the emission handedness arises from collective circulation of light in a twisted bilayer⁴.

A relative twist between two otherwise achiral photonic membranes (Fig. 1a) creates a moiré bilayer where isotropic guided resonances are localized by the pump into counter-rotating collective modes. The resulting broken mirror symmetry introduces non-Hermitian coupling between the modes that lifts their degeneracy and sets the handedness of the emission, resulting in deterministic

lasing of a single chiral vortex state, without any chiral meta-atom, helical geometry or asymmetric pumping.

Extending the reach of moiré photonics—already a powerful tool to control light-matter interactions^{5–9}—this work further shows that, in twisted photonic structures, chirality can arise as a collective property, enabling angular momentum control and chiral light generation⁴.

Remarkably, the lasing mode obtained by Peng et al.⁴ from their twisted moiré metasurface is a *ring-vortex mode*¹⁰ (Fig. 1b) which combines a circulating phase with a polarization structure. It arises collectively from the symmetry and the coupling of two achiral building blocks.

The starting point is a single square-lattice (D_{4h}) metasurface that supports a transverse-electric (TE) band, whose Γ -point mode transforms according to the A_{2g} representation, responsible for the polarization structure of the final lasing mode. Microscopically, each meta-atom in this band behaves predominantly as a vertically oriented magnetic dipole¹¹, giving rise to a symmetry-protected bound state in the continuum (BIC)¹². In an infinite single structure of this kind, this mode is perfectly confined at the Γ -point and does not radiate.

However, the optical gain introduced at a finite pumping spot imposes a spatial confinement that effectively quantizes the guided states near the Γ -point, producing a family of *collective guided resonances*. Owing to the isotropic dispersion of the underlying TE band,




Fig. 1 | Artistic views. **a** Schematic of twisted photonic structure. The system comprises a pair of twisted and bonded metasurface membranes with InGaAsP multiple quantum wells (MQWs) separated by a gap distance of 100 nm (Adapted

from Fig. 1 of ref. 4). **b** Artistic view of the ring-vortex mode at different time instants over one optical period. Electric field polarization is shown with arrows, intensity is shown with color.

these states naturally organize into modes with a well-defined angular phase winding of $\pm 2\pi\ell$, $\ell \in \mathbb{N}$, corresponding to clockwise and counter-clockwise rotating states, respectively. When two such metasurfaces are stacked with a relative twist, the resulting moiré superlattice couples the modes of the upper and lower layers, lifting the degeneracy between the counter-rotating states. Additionally, the moiré periodicity is chosen to make diffraction channels coincide with accidental BICs, reducing out-of-plane radiation losses. In this way, the moiré pattern simultaneously mediates interlayer coupling and engineers the system's radiative landscape.

The combined effects of gain-induced quantization, twisted interlayer coupling and diffraction control give rise to a single lasing mode with a striking structure. Its intensity profile is doughnut-shaped, its phase winds by 2π around the beam axis (corresponding to $\ell = 1$), and its polarization is predominantly azimuthal. Crucially, only one of the two possible azimuthal chiral modes reaches threshold, leading to deterministic selection of the lasing handedness without relying on chiral pumping or structural asymmetry at the meta-atom level.

Beyond demonstrating a vortex laser, the study links collective guided resonances, twist-induced symmetry breaking and non-Hermitian interlayer coupling into a single mechanism that selects a chiral lasing mode. More broadly, it suggests that moiré engineering can be used not only to tailor dispersion but also to control the very topology of optical motion, opening new directions for chiral light sources and collective photonic states.

By tuning and scaling the mechanism reported by Peng and co-workers, for example, by varying twist angle, layer spacing or pumping conditions, twisted bilayer metasurfaces could provide a compact route to on-chip chiral lasers and a broader variety of engineered emission states.

Kristina Frizyuk¹✉ & **Costantino De Angelis**^{2,3}✉

¹Institut für Theoretische Festkörperphysik, Karlsruhe Institute of Technology, Karlsruhe, Germany. ²Department of Information Engineering, University of Brescia, Brescia, Italy. ³National Institute of Optics-National Research Council, Brescia, Italy.

✉ e-mail: kristina.frizyuk@kit.edu; costantino.deangelis@unibs.it

Received: 17 February 2026; Accepted: 26 February 2026;

Published online: 12 March 2026

References

1. Zhang, X., Liu, Y., Han, J., Kivshar, Y. & Song, Q. Chiral emission from resonant metasurfaces. *Science* **377**, 1215–1218 (2022).
2. Deng, H. et al. Chiral lasing enabled by strong coupling. *Sci. Adv.* **11**, eads9562 (2025).
3. Gromyko, D. et al. Unidirectional chiral emission via twisted bi-layer metasurfaces. *Nat. Commun.* **15**, 9804 (2024).
4. Peng, C. et al. Orbital chiral lasing in twisted bilayer metasurfaces. Preprint at: <https://doi.org/10.21203/rs.3.rs-7424101/v1> (2025).
5. Liu, S. et al. Moiré metasurfaces for dynamic beamforming. *Sci. Adv.* **8**, eabo1511 (2022).
6. Tang, H. et al. An adaptive moiré sensor for spectro-polarimetric hyperimaging. *Nat. Photonics* **19**, 463–470 (2025).
7. Du, L. et al. Nonlinear physics of moiré superlattices. *Nat. Mater.* **23**, 1179–1192 (2024).
8. Zhang, T. et al. Twisted moiré photonic crystal enabled optical vortex generation through bound states in the continuum. *Nat. Commun.* **14**, 6014 (2023).
9. Lou, B. et al. Free-space beam steering with twisted bilayer photonic crystal slabs. *ACS Photonics* **11**, 3636–3643 (2024).
10. Chen, Y. et al. Observation of chiral emission enabled by collective guided resonances. *Nat. Nanotechnol.* **20**, 1205–1212 (2025).
11. Sadrieva, Z., Frizyuk, K., Petrov, M., Kivshar, Y. & Bogdanov, A. Multipolar origin of bound states in the continuum. *Phys. Rev. B* **100**, 115303 (2019).
12. Hsu, C. W. et al. Observation of trapped light within the radiation continuum. *Nature* **499**, 188 (2013).

Author contributions

K.F. and C.D.A. contributed equally to all aspects of this work.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Kristina Frizyuk or Costantino De Angelis.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2026